

# Acoustic Radiation Force on a Sphere in Tissue

Yurii A. Ilinskii, Evgenia A. Zabolotskaya and Mark F. Hamilton

*Applied Research Laboratories, The University of Texas at Austin, Austin, Texas 78713–8029*

**Abstract.** A theory is presented for the acoustic radiation force on a sphere embedded in a soft elastic medium that possesses a shear modulus several orders of magnitude smaller than its bulk modulus. Scattering of both compressional and shear waves is taken into account. There is no restriction on the size of the sphere or, apart from symmetry about an axis passing through the center of the sphere, the form of the incident compressional wave. If the medium is a liquid and the sphere is small in comparison with a wavelength the result reduces to the classical theory developed by Gor'kov [1] and others. The effect of shear elasticity on the radiation force is discussed in the context of parameters relevant to a gas bubble in tissue.

**Keywords:** acoustic radiation force, gas bubble, tissue

**PACS:** 43.25.Qp, 43.35.Cg

## INTRODUCTION

Theory for acoustic radiation force on a particle in liquid is well established; see, for example, the review by Wang and Lee [2]. Presented here is a theory for the radiation force on a sphere of arbitrary size, embedded in a soft, homogeneous, isotropic elastic medium. By “soft” is meant a medium characterized by a shear modulus that is several orders of magnitude smaller than the bulk modulus, which is the case for soft tissue. The calculation presented here differs from previous analyses for liquids in that Lagrangian coordinates are used. Care must be taken when calculating the radiation force in Lagrangian coordinates to remove elements of the stress tensor that are associated only with volume potential forces. The effect of shear elasticity on the radiation force is evaluated for parameters corresponding to a gas bubble in tissue. For a small sphere in liquid, classical results obtained by Gor'kov [1] and others are recovered.

## SOLUTION PROCEDURE

The incident and scattered compressional waves, and the scattered shear wave, are defined by the scalar potential functions  $\psi$  and  $\Pi$ , respectively, such that the particle displacement in the elastic medium surrounding the sphere is given by  $\mathbf{u} = \nabla\psi + \nabla \times \nabla \times (\mathbf{r}\Pi)$ , where  $\mathbf{r}$  is position relative to the center of the sphere. The potentials for harmonic waves with angular frequency  $\omega$  are expressed in spherical coordinates as

$$\psi = \frac{1}{2} e^{-i\omega t} \sum_{n=0}^{\infty} a_n [j_n(kr) + A_n h_n(kr)] P_n(\cos \theta) + \text{c.c.} \quad (1)$$

$$\Pi = \frac{1}{2} e^{-i\omega t} \sum_{n=1}^{\infty} a_n B_n h_n(\kappa r) P_n(\cos \theta) + \text{c.c.} \quad (2)$$

where  $j_n$  and  $h_n(\equiv h_n^{(1)})$  are the spherical Bessel and Hankel functions, respectively,  $P_n$  the Legendre polynomials,  $k = \omega/c_l$  and  $\kappa = \omega/c_t$  the wave numbers of the compressional and shear waves having propagation speeds  $c_l$  and  $c_t$ , respectively,  $a_n$  the amplitude coefficients of the incident compressional wave, and  $A_n$  and  $B_n$  the relative amplitude coefficients of the scattered compressional and shear waves, respectively ( $a_n$  has dimensions of length squared, whereas  $A_n$  and  $B_n$  are dimensionless). Shear viscosity, thermal conductivity, and hysteretic behavior that characterize biological media are not considered here. The sphere itself is elastic with similar potentials, not presented here, describing the field inside.

The starting point for the calculation is the momentum equation expressed in Lagrangian coordinates as  $\rho_0 \dot{u}_k = \partial \sigma_{kl} / \partial x_l$ , where the dots indicate time derivatives,  $\rho_0$  is the density of the undeformed medium, and  $\sigma_{kl}$  is the Piola-Kirchhoff pseudostress tensor. In Lagrangian coordinates the surface of the sphere is fixed in the reference frame, and all nonlinearity is thus confined to the stress tensor, which is related to the elastic energy density  $\mathcal{E}$  as  $\sigma_{kl} = \partial \mathcal{E} / \partial (\partial u_k / \partial x_l)$ . The formulation of  $\mathcal{E}$  due to Landau and Lifshitz [3] is employed, in which the second-order elastic constants are the bulk modulus  $K$  and shear modulus  $\mu$ , and the third-order elastic constants are  $A$ ,  $B$ , and  $C$ .

The focus of the present work is on the acoustic radiation force exerted on a sphere in a tissue-like medium. For soft tissue the ratio  $\mu/K$  is in the range of  $10^{-4}$  to  $10^{-6}$ , and measurements reported by Catheline et al. [4] reveal that  $A = O(\mu)$  and  $B = O(K)$ . Under these conditions, with the further recognition that  $K + B = O(\mu)$  [5], the expression for the stress tensor through quadratic order in the particle displacement reduces to

$$\sigma_{kl} = \mu \left( \frac{\partial u_k}{\partial x_l} + \frac{\partial u_l}{\partial x_k} \right) + (K - 2\mu/3) \frac{\partial u_m}{\partial x_m} \delta_{kl} - K \frac{\partial u_m}{\partial x_m} \frac{\partial u_l}{\partial x_k} \quad (3)$$

Omitted from this expression are terms that correspond to the volume potential force produced in the elastic medium by the scattered compressional wave. The volume potential force is not needed in the calculation of the desired surface force on the sphere.

Time averaging eliminates the linear terms, and the remaining term is labeled  $\sigma'_{kl} = -K \langle (\partial u_m / \partial x_m) (\partial u_l / \partial x_k) \rangle$ , where  $\langle \dots \rangle$  represents time averaging. The quantity  $\sigma'_{kl}$  also contains a component associated with the volume force described above, as well as the volume force due to the scattered shear wave. The former is denoted  $\sigma''_{kl}$ , which can be determined and subtracted from  $\sigma'_{kl}$  when calculating the surface force on the sphere. This volume force is  $f_k = \partial \sigma_{kl} / \partial x_l$ , which can be expressed as the gradient of a scalar potential function  $P$  as  $f_k = -\partial P / \partial x_k$ , where  $P = \frac{1}{2} K k^2 [k^2 \psi^2 - (\partial \psi / \partial x_m)^2]$ . With  $\sigma''_{kl} = -\langle P \rangle \delta_{kl}$ , the radiation force in the  $x_k$  direction on a sphere of radius  $R$  is

$$F_k = \int_S (\sigma'_{kl} - \sigma''_{kl}) \frac{x_l}{R} dS \quad (4)$$

where  $S$  is the surface of the sphere. Because of symmetry, only the axial force  $F_z$  is nonzero. Linear theory is used to calculate the quadratic terms in  $\sigma'_{kl}$  and  $\sigma''_{kl}$ .

## RESULTS

It is convenient to express the radiation force as  $F_z = F_{z1} + F_{z2}$ , where  $F_{z1}$  involves the scattering of only compressional waves, whereas  $F_{z2}$  involves the scattering of both compressional and shear waves. Since for liquids  $F_{z2} = 0$ , one may anticipate that  $F_{z1}$

provides the dominant contribution in soft tissue. From Eq. (4) is obtained

$$F_{z1} = i\pi K k^2 \sum_{n=0}^{\infty} \frac{n+1}{(2n+1)(2n+3)} a_n^* a_{n+1} (A_n^* + A_{n+1} + 2A_n^* A_{n+1}) + \text{c.c.} \quad (5)$$

$$F_{z2} = -\pi K k^2 \sum_{n=1}^{\infty} \frac{(\kappa R) h_n(\kappa R)}{2n+1} [j_n(kR) + A_n^* h_n^*(kR)] \\ \times a_n^* \left[ \frac{n(n+1)(n+2)}{2n+3} a_{n+1} B_{n+1} - \frac{(n-1)n(n+1)}{2n-1} a_{n-1} B_{n-1} \right] + \text{c.c.} \quad (6)$$

These expressions are valid for all  $kR$  and  $\kappa R$ , and it may be observed that  $F_{z2}$  vanishes in an oscillatory fashion as  $kR$  and  $\kappa R$  tend toward infinity. The incident wave coefficients  $a_n$  are prescribed and the scattered wave coefficients  $A_n$  and  $B_n$ , which are functions of  $kR$  and  $\kappa R$ , are calculated by solving the linear boundary value problem [6].

Considered first is  $F_{z1}$  for a small sphere ( $kR \ll 1$ ). In this case it is sufficient to retain only the first term in the summation in Eq. (5), such that

$$F_{z1} = a_0^* a_1 (A_0^* + A_1 + 2A_0^* A_1) \frac{i\pi}{3} K k^2 + \text{c.c.} \quad (7)$$

If the incident field is the travelling compressional plane wave  $\psi = \frac{1}{2} \psi_0 e^{i(kz - \omega t)} + \text{c.c.}$ , for example, then  $a_n = i^n (2n+1) \psi_0$ . Regardless of the choice of incident field, for  $kR \ll 1$  the coefficients for the monopole ( $A_0$ ) and dipole ( $A_1$ ) scattering terms are

$$A_0 = \frac{i}{3} (kR)^3 \left( \frac{K}{K_p} - 1 \right) \left[ \left( 1 + \frac{4\mu}{3K_p} \right) - \frac{\omega^2}{\omega_R^2} - \frac{i}{3} (kR)^3 \left( \frac{K}{K_p} - 1 \right) \right]^{-1} \quad (8)$$

$$A_1 = -\frac{i}{3} (kR)^3 \frac{\rho - \rho_p}{\rho + 2\rho_p} \left[ 1 + \frac{i}{3} (kR)^3 \frac{\rho - \rho_p}{\rho + 2\rho_p} \right]^{-1} \quad (9)$$

where  $K_p$  and  $\rho_p$  are the bulk modulus and density of the sphere (particle), respectively, and  $\omega_R = (3K_p/\rho R^2)^{1/2}$  is the natural frequency of the sphere in the absence of shear forces. For  $\mu = 0$ , substitution of the expressions for the incident and scattered wave coefficients in Eq. (7) recovers exactly the result obtained by Gor'kov [1].

To evaluate the effect of finite shear modulus it is helpful to simplify Eq. (8) for the case of a gas bubble ( $K/K_p \gg 1$ ,  $\rho/\rho_p \gg 1$ ) driven below resonance ( $\omega^2/\omega_R^2 \ll 1$ ) and ignore radiation damping, which is associated with the last term inside the square brackets. With  $K_p = \gamma P_0$ , where  $\gamma$  is the ratio of specific heats for the gas and  $P_0$  is atmospheric pressure, Eq. (8) reduces to

$$A_0 \simeq \frac{i}{3} (kR)^3 \frac{K}{\gamma P_0} \left( 1 + \frac{4\mu}{3\gamma P_0} \right)^{-1} \quad (10)$$

Since  $\rho \gg \rho_p$  the magnitude of  $A_0/A_1$  is of order  $K/\gamma P_0 \sim 10^4$ , so the contribution from the dipole term is negligible, and  $F_{z1}$  is thus proportional to  $A_0$ . Reported measurements of shear moduli for soft tissue have values ranging from on the order of 1 kPa on the low end (breast, liver) up to 70 kPa on the high end (cancerous tissue) [7]. Taking  $\mu = 70$  kPa,  $P_0 = 100$  kPa, and  $\gamma = 1.4$  yields  $\mu/\gamma P_0 = 0.5$ , for which  $F_{z1}$  is 40% less than the value predicted for a gas bubble in liquid.

To assess the contribution due to  $F_{z2}$  only the lowest-order scattering of the shear wave is considered here. Since there is no monopole scattering of a shear wave by the sphere ( $B_0 = 0$ ), the lowest-order contribution is dipole scattering ( $B_1$ ). With  $B_n = 0$  for  $n \neq 1$  Eq. (6) becomes

$$F_{z2} = a_1 a_2^* B_1 \frac{2\pi}{5} K k^2 (\kappa R) h_2(\kappa R) [j_2(kR) + A_2^* h_2^*(kR)] + \text{c.c.} \quad (11)$$

At leading order  $F_{z2}$  thus requires participation of the quadrupole component of the incident wave field ( $a_2$ ), whereas the highest-order multipole required by  $F_{z1}$  is the dipole component ( $a_1$ ).

## SUMMARY

The theory presented here provides the radiation force on an isotropic elastic sphere embedded in a soft elastic medium due to a compressional wave incident on the sphere. The incident field is assumed axisymmetric but is otherwise arbitrary, and there is no restriction on the size of the sphere. The force is separated into two components, one that is independent of the scattered shear wave ( $F_{z1}$ ), and another that depends on the scattered shear wave as well as the scattered compressional wave ( $F_{z2}$ ). For small spheres  $F_{z2}$  is negligible, and for  $F_{z1}$  it is sufficient to use Eqs. (7)–(9), which show explicitly the contribution due to shear elasticity in the medium surrounding the particle. For large spheres both  $F_{z1}$  and  $F_{z2}$  contribute to the radiation force, and it is important to note a distinction between the two contributions. The removal of the volume potential force associated with the scattered compressional wave results in an expression for  $F_{z1}$  that is the desired force, i.e., that which produces finite displacement of the sphere. In contrast, the volume force associated with the scattered shear wave is included in  $F_{z2}$ , so while  $F_{z2}$  is the true force acting on a stationary sphere, it is not the force that produces finite displacement of the sphere. One possible avenue for future work is to calculate the influence of the volume force on displacement of the sphere.

## ACKNOWLEDGMENTS

This work was supported by NIH grant numbers DK070618 and EB011603.

## REFERENCES

1. L. P. Gor'kov, "On the forces acting on a small particle in an acoustical field in an ideal fluid," *Sov. Phys. Doklady* **6**, 773–775 (1962).
2. T. G. Wang and C. P. Lee, "Radiation pressure and acoustic levitation," Chap. 6 in *Nonlinear Acoustics*, M. F. Hamilton and D. T. Blackstock, eds. (Acoust. Soc. Am., New York, 2008), pp. 177–205.
3. L. D. Landau and E. M. Lifshitz, *Theory of Elasticity*, 3rd ed. (Pergamon Press, Oxford, 1986), p. 107.
4. S. Catheline, J.-L. Gennisson, and M. Fink, "Measurement of elastic nonlinearity of soft solid with transient elastography," *J. Acoust. Soc. Am.* **114**, 3087–3091 (2003).
5. M. F. Hamilton, Yu. A. Ilinskii, and E. A. Zabolotskaya, "Separation of compressibility and shear deformation in the elastic energy density," *J. Acoust. Soc. Am.* **116**, 41–44 (2004).
6. C. F. Ying and R. Truell, "Scattering of a plane longitudinal wave by a spherical obstacle in an isotropically elastic solid," *J. Appl. Phys.* **27**, 1086–1097 (1956).
7. P. N. T. Wells and H.-D. Liang, "Medical ultrasound: imaging of soft tissue strain and elasticity," *J. R. Soc. Interface* **8**, 1521–1549 (2011).

Copyright of AIP Conference Proceedings is the property of American Institute of Physics and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.