

4-[Tris(1*H*-pyrazol-1-yl)methyl]phenol

Xiao-Yan Chen, Xiaoping Yang and Bradley J. Holliday*

 Department of Chemistry and Biochemistry, University of Texas at Austin, 1 University Station, A5300, Austin, TX 78712, USA
 Correspondence e-mail: bholliday@cm.utexas.edu


Received 9 August 2011; accepted 17 October 2011

 Key indicators: single-crystal X-ray study; $T = 153$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.043; wR factor = 0.099; data-to-parameter ratio = 12.5.

The title compound, $\text{C}_{16}\text{H}_{14}\text{N}_6\text{O}$, was prepared by the condensation of 4-(trifluoromethyl)phenol and sodium pyrazol-1-ide in a yield of 58%. The dihedral angles formed by the planes of the pyrazole rings are 50.7 (2), 71.2 (3) and 95.8 (2)°. The molecules are associated into dimers by pairs of intermolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bonds involving the hydroxy groups and pyrazole N atoms. In addition, $\pi-\pi$ stacking between the phenol rings of these inversion-related dimers is observed, with a ring centroid-to-centroid distance of 3.9247 (10) Å.

Related literature

For the preparation and coordination chemistry of tris(pyrazolyl)borates and tris(pyrazolyl)methanes, see: Trofimenko (1966, 1970, 1999); Pettinari & Pettinari (2005); Reger *et al.* (2000). For the chemistry of tris(pyrazolyl)methane derivatives, see: Humphrey *et al.* (1999). For similar structures, see: Liddle & Gardinier (2007).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{14}\text{N}_6\text{O}$	$\gamma = 109.40$ (3)°
$M_r = 306.33$	$V = 746.0$ (3) Å ³
Triclinic, $P\bar{1}$	$Z = 2$
$a = 8.5065$ (17) Å	Mo $K\alpha$ radiation
$b = 8.6829$ (17) Å	$\mu = 0.09$ mm ⁻¹
$c = 10.815$ (2) Å	$T = 153$ K
$\alpha = 96.97$ (3)°	$0.30 \times 0.28 \times 0.20$ mm
$\beta = 91.51$ (2)°	

Data collection

Nonius KappaCCD diffractometer	4225 measured reflections
Absorption correction: Gaussian (XPRED in SHELXTL; Sheldrick, 2008)	2609 independent reflections
$T_{\min} = 0.973$, $T_{\max} = 0.983$	1975 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$	209 parameters
$wR(F^2) = 0.099$	H-atom parameters constrained
$S = 1.07$	$\Delta\rho_{\text{max}} = 0.18$ e Å ⁻³
2609 reflections	$\Delta\rho_{\text{min}} = -0.27$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1A}\cdots\text{N4}^i$	0.82	2.02	2.836 (2)	173

 Symmetry code: (i) $-x + 2, -y + 2, -z$.

Data collection: COLLECT (Nonius, 1998); cell refinement: SCALEPACK (Otwinowski & Minor, 1997); data reduction: DENZO-SMN (Otwinowski & Minor, 1997); program(s) used to solve structure: SIR97 (Altomare *et al.*, 1999); program(s) used to refine structure: XL in SHELXTL/PC (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and POV-RAY (Persistence of Vision Team, 2004); software used to prepare material for publication: SHELXL97 (Sheldrick, 2008) and WinGX (Farrugia, 1999).

The authors gratefully acknowledge the Robert A. Welch Foundation (grant No. F-1631), the National Science Foundation (grant Nos. CHE-0741973 and CHE-0847763), the Advanced Research Program of the Texas Higher Education Coordinating Board (grant No. 01916-090-2010) and the University of Texas at Austin for financial support of this research.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2340).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Humphrey, E. R., Mann, K. L. V., Reeves, Z. R., Behrendt, A., Jeffery, J. C., Maher, J. P., McCleverty, J. A. & Ward, M. D. (1999). *New J. Chem.* **23**, 417–423.
- Liddle, B. & Gardinier, J. R. (2007). *J. Org. Chem.* **72**, 9794–9797.
- Nonius (1998). COLLECT. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Persistence of Vision Team (2004). POV-RAY. Persistence of Vision Raytracer Pty. Ltd, Victoria, Australia. <http://www.povray.org/>
- Pettinari, C. & Pettinari, R. (2005). *Coord. Chem. Rev.* **249**, 525–543.
- Reger, D. L., Grattan, T. C., Brown, K. J., Little, C. A., Lamba, J. J. S., Rheingold, A. L. & Sommer, R. D. (2000). *J. Organomet. Chem.* **607**, 120–128.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Trofimenko, S. (1966). *J. Am. Chem. Soc.* **88**, 1842–1844.
- Trofimenko, S. (1970). *J. Am. Chem. Soc.* **92**, 5118–5126.
- Trofimenko, S. (1999). *Scorpionates: The Coordination Chemistry of Polypyrazolylborate Ligands*. London: Imperial College Press.

supplementary materials

Acta Cryst. (2011). E67, o3045 [doi:10.1107/S1600536811043042]

4-[Tris(1*H*-pyrazol-1-yl)methyl]phenol

X.-Y. Chen, X. Yang and B. J. Holliday

Comment

Since seminal reports on tris(pyrazolyl)borates and tris(pyrazolyl)methanes (Trofimenko, 1966, 1970), variations on these ligands have been widely studied. Currently, there are great research efforts to study ligands of this type that are functionalized at the back boron or methine positions. The synthesis and coordination chemistry of aryltris(pyrazolyl)borates have also been well investigated, which has allowed significant advances in iron(II) spin-crossover chemistry. However, the chemistry of analogous tris(pyrazolyl)methane derivatives is almost unknown. Herein, we demonstrate the preparation of a new tris(pyrazolyl)methane derivative, 4-[tris(1*H*-pyrazol-1-yl)methyl]phenol, using a one-pot synthesis method with 4-(trifluoromethyl)phenol and freshly prepared sodium pyrazol-1-ide to give the desired product in good yield.

The solid state structure of 4-[tris(1*H*-pyrazol-1-yl)methyl]phenol can be seen in Fig. 1. There are two molecules in the unit cell. The molecules are associated into dimers by pairs of intermolecular O—H···N hydrogen bonds involving the hydroxyl groups and pyrazole N atoms. In addition, π – π stacking between the phenol rings of these inversion-related ($-x + 2, -y + 2, -z$) dimers is observed with a ring centroid-to-centroid distance of 3.9247 (10) Å.

Experimental

The title compound was prepared by refluxing 4-(trifluoromethyl)phenol (0.162 g, 1.0 mmol) and freshly prepared sodium pyrazol-1-ide (0.361 g, 4.0 mmol) in tetrahydrofuran (20 ml) for 12 h under a nitrogen atmosphere. The desired product was purified by column chromatography on silica gel using CH₂Cl₂ as the eluent with a yield of 58%. Single crystals suitable for X-ray diffraction were obtained *via* slow evaporation from a methanol solution.

Refinement

All H atoms were positioned geometrically and refined using a riding model, with C—H = 0.95 Å, and O—H = 0.82 Å. $U_{\text{iso}}(\text{H})$ values were set to $1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. Ellipsoid plot of C₁₆H₁₄N₆O showing selected atoms at 30% probability level.




Fig. 2. A packing diagram of $C_{16}H_{14}N_6O$.

4-[Tris(1*H*-pyrazol-1-yl)methyl]phenol

Crystal data

$C_{16}H_{14}N_6O$	$Z = 2$
$M_r = 306.33$	$F(000) = 320$
Triclinic, $P\bar{1}$	$D_x = 1.364 \text{ Mg m}^{-3}$
Hall symbol: $-P 1$	Melting point: 443 K
$a = 8.5065 (17) \text{ \AA}$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$b = 8.6829 (17) \text{ \AA}$	Cell parameters from 2208 reflections
$c = 10.815 (2) \text{ \AA}$	$\theta = 2.9\text{--}27.5^\circ$
$\alpha = 96.97 (3)^\circ$	$\mu = 0.09 \text{ mm}^{-1}$
$\beta = 91.51 (2)^\circ$	$T = 153 \text{ K}$
$\gamma = 109.40 (3)^\circ$	Block, colourless
$V = 746.0 (3) \text{ \AA}^3$	$0.30 \times 0.28 \times 0.20 \text{ mm}$

Data collection

Nonius KappaCCD diffractometer	2609 independent reflections
Radiation source: fine-focus sealed tube graphite	1975 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.025$
Absorption correction: gaussian (<i>XPRED</i> in <i>SHELXTL</i> ; Sheldrick, 2008)	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.9^\circ$
$T_{\text{min}} = 0.973$, $T_{\text{max}} = 0.983$	$h = -10 \rightarrow 10$
4225 measured reflections	$k = -10 \rightarrow 10$
	$l = -12 \rightarrow 11$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.043$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.099$	H-atom parameters constrained
$S = 1.07$	$w = 1/[\sigma^2(F_o^2) + (0.038P)^2 + 0.1787P]$
2609 reflections	where $P = (F_o^2 + 2F_c^2)/3$
209 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.27 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	1.00800 (16)	1.32161 (15)	0.03852 (13)	0.0266 (3)
H1A	0.9790	1.2970	-0.0362	0.054 (8)*
N1	0.74972 (17)	0.78542 (18)	0.40566 (13)	0.0180 (4)
N2	0.69159 (19)	0.65739 (19)	0.47332 (15)	0.0244 (4)
N3	0.98104 (17)	0.72693 (18)	0.31464 (14)	0.0181 (4)
N4	1.08607 (18)	0.73336 (19)	0.22092 (15)	0.0237 (4)
N5	0.70790 (17)	0.60541 (17)	0.21695 (14)	0.0173 (4)
N6	0.54566 (18)	0.59411 (18)	0.19620 (14)	0.0216 (4)
C1	0.9528 (2)	1.1846 (2)	0.09625 (17)	0.0197 (4)
C2	0.8251 (2)	1.0424 (2)	0.04486 (17)	0.0213 (4)
H2A	0.7695	1.0394	-0.0311	0.026*
C3	0.7802 (2)	0.9051 (2)	0.10623 (16)	0.0191 (4)
H3A	0.6944	0.8103	0.0711	0.023*
C4	0.8613 (2)	0.9069 (2)	0.21937 (16)	0.0165 (4)
C5	0.9873 (2)	1.0523 (2)	0.27129 (17)	0.0206 (4)
H5A	1.0427	1.0560	0.3475	0.025*
C6	1.0305 (2)	1.1896 (2)	0.21160 (17)	0.0213 (4)
H6A	1.1121	1.2863	0.2487	0.026*
C7	0.8240 (2)	0.7575 (2)	0.28833 (16)	0.0173 (4)
C8	0.6124 (2)	0.7133 (2)	0.56240 (18)	0.0250 (5)
H8A	0.5599	0.6528	0.6241	0.030*
C9	0.6169 (2)	0.8729 (2)	0.55301 (18)	0.0263 (5)
H9A	0.5710	0.9371	0.6055	0.032*
C10	0.7026 (2)	0.9154 (2)	0.45071 (17)	0.0208 (4)
H10A	0.7246	1.0141	0.4180	0.025*
C11	1.2086 (2)	0.6894 (2)	0.2675 (2)	0.0268 (5)
H11A	1.2999	0.6832	0.2244	0.032*
C12	1.1831 (2)	0.6538 (2)	0.3889 (2)	0.0281 (5)
H12A	1.2517	0.6210	0.4405	0.034*
C13	1.0367 (2)	0.6771 (2)	0.41634 (18)	0.0225 (5)
H13A	0.9846	0.6617	0.4906	0.027*
C14	0.4724 (2)	0.4422 (2)	0.13858 (17)	0.0220 (4)
H14A	0.3599	0.3985	0.1111	0.026*

supplementary materials

C15	0.5835 (2)	0.3552 (2)	0.12394 (19)	0.0284 (5)
H15A	0.5602	0.2468	0.0870	0.034*
C16	0.7341 (2)	0.4636 (2)	0.17562 (18)	0.0238 (5)
H16A	0.8347	0.4434	0.1811	0.029*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0374 (8)	0.0219 (8)	0.0236 (9)	0.0116 (6)	0.0085 (6)	0.0092 (6)
N1	0.0197 (8)	0.0170 (8)	0.0171 (8)	0.0049 (7)	0.0029 (7)	0.0049 (7)
N2	0.0284 (9)	0.0226 (9)	0.0218 (9)	0.0056 (7)	0.0055 (7)	0.0093 (7)
N3	0.0155 (8)	0.0185 (9)	0.0209 (9)	0.0065 (6)	0.0013 (7)	0.0034 (7)
N4	0.0187 (8)	0.0256 (9)	0.0297 (10)	0.0101 (7)	0.0072 (7)	0.0066 (8)
N5	0.0149 (8)	0.0167 (9)	0.0207 (9)	0.0063 (6)	0.0007 (6)	0.0017 (7)
N6	0.0152 (8)	0.0231 (9)	0.0255 (9)	0.0057 (7)	0.0005 (7)	0.0023 (7)
C1	0.0231 (10)	0.0180 (11)	0.0234 (11)	0.0126 (8)	0.0098 (8)	0.0056 (8)
C2	0.0241 (10)	0.0261 (11)	0.0178 (10)	0.0133 (9)	0.0009 (8)	0.0044 (9)
C3	0.0184 (10)	0.0202 (11)	0.0183 (10)	0.0067 (8)	0.0003 (8)	0.0001 (8)
C4	0.0156 (9)	0.0167 (10)	0.0189 (10)	0.0077 (8)	0.0041 (8)	0.0025 (8)
C5	0.0181 (10)	0.0235 (11)	0.0199 (11)	0.0070 (8)	-0.0014 (8)	0.0020 (8)
C6	0.0210 (10)	0.0167 (10)	0.0235 (11)	0.0032 (8)	0.0030 (8)	0.0011 (8)
C7	0.0145 (9)	0.0186 (10)	0.0188 (10)	0.0061 (8)	0.0008 (8)	0.0017 (8)
C8	0.0210 (10)	0.0329 (12)	0.0185 (11)	0.0049 (9)	0.0037 (8)	0.0048 (9)
C9	0.0228 (11)	0.0321 (12)	0.0248 (11)	0.0120 (9)	0.0039 (9)	-0.0010 (9)
C10	0.0206 (10)	0.0201 (10)	0.0241 (11)	0.0106 (8)	-0.0002 (8)	0.0013 (8)
C11	0.0181 (10)	0.0227 (11)	0.0415 (14)	0.0090 (9)	0.0027 (9)	0.0056 (10)
C12	0.0228 (11)	0.0230 (11)	0.0400 (14)	0.0101 (9)	-0.0097 (9)	0.0057 (10)
C13	0.0255 (11)	0.0191 (11)	0.0223 (11)	0.0066 (8)	-0.0046 (8)	0.0046 (8)
C14	0.0190 (10)	0.0220 (11)	0.0199 (11)	0.0005 (8)	0.0004 (8)	0.0023 (8)
C15	0.0305 (12)	0.0193 (11)	0.0309 (12)	0.0053 (9)	0.0004 (9)	-0.0042 (9)
C16	0.0253 (11)	0.0198 (11)	0.0297 (12)	0.0128 (9)	0.0023 (9)	0.0012 (9)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.360 (2)	C4—C5	1.399 (3)
O1—H1A	0.8200	C4—C7	1.521 (3)
N1—C10	1.360 (2)	C5—C6	1.371 (3)
N1—N2	1.366 (2)	C5—H5A	0.9300
N1—C7	1.462 (2)	C6—H6A	0.9300
N2—C8	1.325 (2)	C8—C9	1.390 (3)
N3—C13	1.358 (2)	C8—H8A	0.9300
N3—N4	1.363 (2)	C9—C10	1.362 (3)
N3—C7	1.473 (2)	C9—H9A	0.9300
N4—C11	1.330 (2)	C10—H10A	0.9300
N5—C16	1.349 (2)	C11—C12	1.391 (3)
N5—N6	1.361 (2)	C11—H11A	0.9300
N5—C7	1.470 (2)	C12—C13	1.361 (3)
N6—C14	1.323 (2)	C12—H12A	0.9300
C1—C6	1.385 (3)	C13—H13A	0.9300

C1—C2	1.387 (3)	C14—C15	1.393 (3)
C2—C3	1.382 (3)	C14—H14A	0.9300
C2—H2A	0.9300	C15—C16	1.369 (3)
C3—C4	1.385 (2)	C15—H15A	0.9300
C3—H3A	0.9300	C16—H16A	0.9300
C1—O1—H1A	109.5	N1—C7—N3	109.47 (14)
C10—N1—N2	111.56 (15)	N5—C7—N3	107.01 (14)
C10—N1—C7	128.86 (15)	N1—C7—C4	110.73 (14)
N2—N1—C7	118.61 (14)	N5—C7—C4	113.65 (14)
C8—N2—N1	104.13 (15)	N3—C7—C4	108.88 (14)
C13—N3—N4	111.24 (14)	N2—C8—C9	112.15 (18)
C13—N3—C7	130.58 (16)	N2—C8—H8A	123.9
N4—N3—C7	117.89 (14)	C9—C8—H8A	123.9
C11—N4—N3	104.65 (15)	C10—C9—C8	105.46 (18)
C16—N5—N6	112.33 (15)	C10—C9—H9A	127.3
C16—N5—C7	129.11 (15)	C8—C9—H9A	127.3
N6—N5—C7	118.30 (14)	N1—C10—C9	106.65 (17)
C14—N6—N5	103.95 (14)	N1—C10—H10A	126.7
O1—C1—C6	117.37 (17)	C9—C10—H10A	126.7
O1—C1—C2	123.26 (17)	N4—C11—C12	111.55 (17)
C6—C1—C2	119.36 (17)	N4—C11—H11A	124.2
C3—C2—C1	120.19 (17)	C12—C11—H11A	124.2
C3—C2—H2A	119.9	C13—C12—C11	105.54 (17)
C1—C2—H2A	119.9	C13—C12—H12A	127.2
C2—C3—C4	120.83 (17)	C11—C12—H12A	127.2
C2—C3—H3A	119.6	N3—C13—C12	107.00 (18)
C4—C3—H3A	119.6	N3—C13—H13A	126.5
C3—C4—C5	118.31 (17)	C12—C13—H13A	126.5
C3—C4—C7	123.42 (16)	N6—C14—C15	112.14 (17)
C5—C4—C7	118.25 (16)	N6—C14—H14A	123.9
C6—C5—C4	121.01 (17)	C15—C14—H14A	123.9
C6—C5—H5A	119.5	C16—C15—C14	105.15 (17)
C4—C5—H5A	119.5	C16—C15—H15A	127.4
C5—C6—C1	120.22 (17)	C14—C15—H15A	127.4
C5—C6—H6A	119.9	N5—C16—C15	106.41 (17)
C1—C6—H6A	119.9	N5—C16—H16A	126.8
N1—C7—N5	106.98 (14)	C15—C16—H16A	126.8

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1A \cdots N4 ⁱ	0.82	2.02	2.836 (2)	173.

Symmetry codes: (i) $-x+2, -y+2, -z$.

Fig. 1




Fig. 2

