

Copyright  
by  
Elaine Marie Cozzi  
2007

The Dissertation Committee for Elaine Marie Cozzi  
certifies that this is the approved version of the following dissertation:

## **Incompressible Fluids with Vorticity in Besov Spaces**

Committee:

---

Mikhail M. Vishik, Supervisor

---

William Beckner

---

Rafael de la Llave

---

Hans Koch

---

Philip J. Morrison

**Incompressible Fluids with Vorticity in Besov Spaces**

by

**Elaine Marie Cozzi, B.A.**

**DISSERTATION**

Presented to the Faculty of the Graduate School of  
The University of Texas at Austin  
in Partial Fulfillment  
of the Requirements  
for the Degree of

**DOCTOR OF PHILOSOPHY**

THE UNIVERSITY OF TEXAS AT AUSTIN

August 2007

## Acknowledgments

First and foremost, I wish to thank my parents for teaching me the value of knowledge and a good education. I also thank them for working day and night so that I may have the opportunity to go to graduate school and pursue a degree in mathematics.

There are two people who played a vital role in my production of this thesis. The first is my advisor, Misha Vishik, who convinced me that I was capable of solving difficult open problems at a time when I did not believe this was possible. I especially thank him for always making himself available to offer advice, encouragement, and mathematical insight, without ever being overbearing. Finally, I thank him for his patience and his sense of humor during our weekly meetings.

The second person who played a fundamental role in my growth as a mathematician is my academic older brother, Jim Kelliher. I thank Jim for being there to talk about mathematics any day or night of the week, for offering advice and encouragement, and for just being the most supportive academic big brother anybody could ask for.

I thank Magda Czubak for being my friend, for making me laugh, and for making late nights at RLM oddly enjoyable. I also thank Magda for inspiring me to work harder through her own passion and determination.

Finally, I thank Clay Petsche for teaching me that I should not feel guilty for taking the day off, and for informing me in my early years of graduate school that there are more important things than preliminary exams. This realization helped carry me through some very challenging and frustrating years.

# Incompressible Fluids with Vorticity in Besov Spaces

Publication No. \_\_\_\_\_

Elaine Marie Cozzi, Ph.D.  
The University of Texas at Austin, 2007

Supervisor: Mikhail M. Vishik

In this thesis, we consider solutions to the two-dimensional Euler equations with uniformly continuous initial vorticity in a critical or subcritical Besov space. We use paradifferential calculus to show that the solution will lose an arbitrarily small amount of smoothness over any fixed finite time interval. This result is motivated by a theorem of Bahouri and Chemin which states that the Sobolev exponent of a solution to the two-dimensional Euler equations in a critical or subcritical Sobolev space may decay exponentially with time. To prove our result, one can use methods similar to those used by Bahouri and Chemin for initial vorticity in a Besov space with Besov exponent between 0 and 1; however, we use different methods to prove a result which applies for any Sobolev exponent between 0 and 2.

The remainder of this thesis is based on joint work with J. Kelliher. We study the vanishing viscosity limit of solutions of the Navier-Stokes equations to solutions of the Euler equations in the plane assuming initial vorticity is in

a variant Besov space introduced by Vishik. Our methods allow us to extend a global in time uniqueness result established by Vishik for the two-dimensional Euler equations in this space.

# Table of Contents

<b>Acknowledgments</b>	<b>iv</b>
<b>Abstract</b>	<b>vi</b>
<b>Chapter 1. Introduction</b>	<b>1</b>
1.1 Chapter summaries . . . . .	2
1.2 Notational conventions . . . . .	5
<b>Chapter 2. Background</b>	<b>6</b>
2.1 Littlewood-Paley theory . . . . .	6
2.2 Function spaces . . . . .	8
2.3 Properties of ideal incompressible fluids . . . . .	11
2.3.1 The Biot-Savart law . . . . .	12
2.3.2 The vorticity of an ideal incompressible fluid . . . . .	13
2.3.3 A few technical lemmas . . . . .	19
<b>Chapter 3. An initial value problem for the Euler equations in the plane</b>	<b>28</b>
3.1 Introduction and history . . . . .	28
3.2 A proof of Theorem 3.1.4 . . . . .	33
3.3 Paradifferential estimates for the vorticity equation . . . . .	38
3.4 Statement and proof of the main results . . . . .	48
3.4.1 Besov regularity of the vorticity . . . . .	48
3.4.2 Besov regularity of the flow . . . . .	52
<b>Chapter 4. Vanishing viscosity in the plane</b>	<b>55</b>
4.1 Introduction and history . . . . .	55
4.2 Statement of the main results . . . . .	56
4.2.1 A comparison of rates of convergence . . . . .	61

4.3	Proof of the main theorem . . . . .	61
4.3.1	Convergence in Equation (4.3.5) . . . . .	64
	<b>Bibliography</b>	<b>68</b>
	<b>Vita</b>	<b>72</b>

# Chapter 1

## Introduction

We begin by introducing the Navier-Stokes equations:

$$(NS) \quad \begin{cases} \partial_t v + v \cdot \nabla v - \nu \Delta v = -\nabla p, \\ \operatorname{div} v = 0, \\ v|_{t=0} = v^0. \end{cases}$$

Here  $v : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  denotes the velocity vector field, and  $p : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$  denotes the pressure. The Navier-Stokes equations model incompressible fluid flow in  $\mathbb{R}^n$ ,  $n \geq 2$ , with constant density and constant viscosity, denoted by  $\nu$ .

When  $\nu = 0$ , the Navier-Stokes equations reduce to the Euler equations:

$$(E) \quad \begin{cases} \partial_t v + v \cdot \nabla v = -\nabla p, \\ \operatorname{div} v = 0, \\ v|_{t=0} = v^0. \end{cases}$$

In this thesis, we study properties of solutions to  $(NS)$  and  $(E)$  in the plane. Specifically, we investigate regularity of solutions to  $(E)$  in the plane and the limit of solutions of  $(NS)$  to solutions of  $(E)$  in the energy norm as  $\nu$  goes to zero.

It is known that when initial data  $v^0$  for  $(E)$  is in the supercritical Sobolev space  $W^{s+1,p}(\mathbb{R}^2)$ , where  $sp > 2$ , the solution does not lose any regularity as time evolves (see, for example, [11], [12]). Much less is known when initial data belongs to the critical Sobolev spaces, where  $sp = 2$ , or to the subcritical Sobolev spaces, where  $sp < 2$ . We therefore restrict our attention to

these two cases. In addition, we assume that the initial vorticity is uniformly continuous.

We also study the vanishing viscosity limit of solutions of the Navier-Stokes equations to solutions of the Euler equations in the plane assuming initial vorticity is in a variant Besov space introduced by Vishik in [21]. Our methods allow us to extend a uniqueness result proved by Vishik.

In the chapters that follow, we begin with an introduction to known results in each of the above areas. We then establish our results. Before we begin, we give a brief summary of each chapter.

## 1.1 Chapter summaries

**Chapter 2:** In this chapter we discuss tools which we utilize throughout Chapters 3 and 4. In particular, we discuss Littlewood-Paley theory and Bony's paraproduct decomposition. We also define useful function spaces. Finally, we state some properties of incompressible fluids in the plane.

**Chapter 3:** In this chapter we study regularity of solutions to  $(E)$  with initial data in a critical or subcritical Sobolev space.

Let  $v = (v_1, v_2)$  be a solution to the two-dimensional Euler equations with vorticity  $\omega(v) = \partial_1 v_2 - \partial_2 v_1$ . Assume  $v$  satisfies  $\omega(v^0) = \omega^0 \in W^{s,p}(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ , where  $sp \leq 2$  and  $s \in (0, 1]$ . Under these assumptions, Bahouri and Chemin find in [1] a lower bound for the Sobolev exponent of  $\omega(t)$  which decays exponentially with time. They show that for all  $\tilde{s} < s$ ,  $\omega(t)$  belongs

to  $W^{\bar{s}\alpha(t),p}(\mathbb{R}^2)$ , where  $\alpha(t) = \exp(-\int_0^t \|v(\tau)\|_{LL} d\tau)$  is the Holder exponent of the flow  $g(t)$ , and  $LL$  denotes the space of log-Lipschitz functions. The authors find this lower bound by studying what happens to the initial vorticity under composition with  $g(t)$ .

Now let  $v$  be the unique solution to the two-dimensional Euler equations such that  $\omega(v^0) = \omega^0 \in UC(\mathbb{R}^2) \cap W^{s,p}(\mathbb{R}^2)$ , for  $sp \leq 2$  with  $p \in (1, \infty)$  and  $s \in (0, 2)$ , where  $UC(\mathbb{R}^2)$  denotes the space of uniformly continuous functions on  $\mathbb{R}^2$ . We show that under these assumptions, the Holder exponent of  $g(t)$  can be made arbitrarily close to 1 over any fixed finite time interval. After a slight modification of the proof in [1], we are able to show that when  $\omega^0 \in UC(\mathbb{R}^2) \cap W^{s,p}(\mathbb{R}^2)$  and  $s \in (0, 1]$ ,  $\omega(t)$  loses an arbitrarily small amount of smoothness in finite time.

Since the methods used by Bahouri and Chemin only apply to the case  $s \in (0, 1]$ , the modification of their proof fails for the case  $s \in (1, 2)$ . To handle this case, we construct a different proof which shows an arbitrarily small loss in smoothness in the appropriate critical and subcritical Besov spaces for *all*  $s \in (0, 2)$ . By embedding properties, one can conclude from this result an arbitrarily small loss of smoothness in Sobolev spaces when vorticity is uniformly continuous.

To prove the result, we consider the localized vorticity equation given by

$$\begin{aligned} \partial_t \Delta_q \omega + v \cdot \nabla \Delta_q \omega &= [v \cdot \nabla, \Delta_q] \omega, \\ \Delta_q \omega|_{t=0} &= \Delta_q \omega^0, \end{aligned} \tag{1.1.1}$$

where  $\Delta_q$  is the Littlewood-Paley operator which projects in Fourier space onto an annulus with inner and outer radii of order  $2^q$ . We then use Bony's paraproduct decomposition given in [3] to bound the commutator on the right hand side of (1.1.1). This estimate, combined with a Gronwall type of argument, implies the result. As  $h(t, x) = g(t)^{-1}(x) - x$  satisfies a similar equation, given by

$$\begin{aligned} \partial_t \Delta_q h + v \cdot \nabla \Delta_q h &= -\Delta_q v + [v \cdot \nabla, \Delta_q] h, \\ \Delta_q h|_{t=0} &= 0, \end{aligned} \tag{1.1.2}$$

we apply similar techniques, combined with the regularity result for the velocity  $v$ , to prove an analogous result for the flow. Specifically, we show that  $h$  belongs to  $L_{loc}^\infty(\mathbb{R}^+; W_p^{s+1-\delta}(\mathbb{R}^2))$  for any fixed  $\delta > 0$ .

**Chapter 4:** The content of this chapter stems from joint work with J. Kelliher. We consider a uniqueness class established by Vishik in [21]. This class is a variant of a critical Besov space. Let  $\Gamma : \mathbb{R} \rightarrow [1, \infty)$  be a locally Lipschitz continuous nondecreasing function with  $\Gamma = 1$  on the interval  $(-\infty, -1]$  and  $\lim_{N \rightarrow \infty} \Gamma(N) = \infty$ . ( $\Gamma$  also satisfies two minor technical conditions which can be found on p. 771 of [21].) Define  $B_\Gamma(\mathbb{R}^2) = \{f \in S'(\mathbb{R}^2) : \sum_{j=-1}^N \|\Delta_j f\|_{L^\infty} \leq C(\Gamma(N))\}$ . In [21], Vishik shows that a solution  $v$  to the two-dimensional Euler equations with  $\omega(v) \in L^\infty([0, T]; L^{p_0}(\mathbb{R}^2) \cap B_\Gamma(\mathbb{R}^2))$  for  $p_0 < 2$  is unique if  $\Gamma(N)$  grows no faster than  $N \log N$ . His result implies, among many things, that if one can construct a solution to the Euler equations in the space  $bmo(\mathbb{R}^2)$ , then the solution must be unique. He proves global

in time existence of solutions in his uniqueness class when initial vorticity belongs to  $B_\Gamma(\mathbb{R}^2) \cap L^{p_0}(\mathbb{R}^2) \cap L^{p_1}(\mathbb{R}^2)$ , where  $\Gamma(N) = \log^\kappa N$  with  $0 < \kappa \leq \frac{1}{2}$ , and  $p_0 < 2 < p_1$ ; he proves short time existence of solutions in his uniqueness class when  $\frac{1}{2} < \kappa \leq 1$ .

We prove that there exists a unique global in time solution  $v$  to the Euler equations in  $L^\infty([0, \infty); H^1(\mathbb{R}^2))$  when initial vorticity belongs to  $B_\Gamma(\mathbb{R}^2) \cap L^2(\mathbb{R}^2)$ , and  $\Gamma(N) = \log^\kappa N$ ,  $0 < \kappa < 1$ . We prove this result *without* showing that the vorticity remains in Vishik's uniqueness class globally in time. We then prove that with this initial data the solutions of the Navier-Stokes equations converge to the unique solution of the Euler equations in the energy norm as viscosity tends to 0.

## 1.2 Notational conventions

We use  $C$  to denote a constant which may differ in value on two sides of an inequality. The constant  $C$  may or may not be absolute; for example,  $C$  will sometimes depend on the initial data. We only clarify this dependence when it is important to the discussion. To demonstrate the dependence of  $C$  on some variable, we often write  $C(*)$ , where  $*$  denotes the variable. If necessary, we distinguish between two unequal constants by writing  $C_1$  and  $C_2$  or  $C$  and  $\tilde{C}$ .

# Chapter 2

## Background

### 2.1 Littlewood-Paley theory

We begin by defining the Littlewood-Paley operators. These operators will play an important role in the proofs of the main results. We start with the following lemma:

**Lemma 2.1.1.** *There exist two radial functions  $\chi \in S(\mathbb{R}^2)$  and  $\varphi \in S(\mathbb{R}^2)$  satisfying the following properties:*

- (i)  $\text{supp } \chi \subset \{\xi \in \mathbb{R}^2 : 0 \leq |\xi| \leq \frac{4}{3}\},$
- (ii)  $\text{supp } \varphi \subset \{\xi \in \mathbb{R}^2 : \frac{3}{4} \leq |\xi| \leq \frac{8}{3}\},$
- (iii)  $\chi(\xi) + \sum_{j=0}^{\infty} \varphi_j(\xi) = 1,$

where  $\varphi_j(\xi) = \varphi(2^{-j}\xi)$  (so  $\check{\varphi}_j(x) = 2^{jn}\check{\varphi}(2^jx)$ ).

*Proof.* This is classical. See for example [19]. □

Observe that, if  $|j - j'| \geq 2$ , then  $\text{supp } \varphi_j \cap \text{supp } \varphi_{j'} = \emptyset$ , and, if  $j \geq 1$ , then  $\text{supp } \varphi_j \cap \text{supp } \chi = \emptyset$ .

Let  $f \in S'(\mathbb{R}^2)$ . We define, for any integer  $j$ ,

$$\Delta_j f = \begin{cases} 0, & j < -1, \\ \chi(D)f = \check{\chi} * f, & j = -1, \\ \varphi(D)f = \check{\varphi}_j * f, & j > -1, \end{cases}$$

and

$$S_j f = \sum_{k=-\infty}^{j-1} \Delta_k f = \chi(2^{-j}D)f.$$

An important tool in chapter 3 will be a decomposition introduced by J.-M. Bony in [3]. We recall the definition of the paraproduct and remainder used in this decomposition.

**Definition 2.1.1.** Define the paraproduct of two functions  $f$  and  $g$  by

$$T_f g = \sum_{\substack{i,j \\ i \leq j-2}} \Delta_i f \Delta_j g = \sum_{j=1}^{\infty} S_{j-1} f \Delta_j g.$$

We use  $R(f, g)$  to denote the remainder.  $R(f, g)$  is given by the following bilinear operator:

$$R(f, g) = \sum_{\substack{i,j \\ |i-j| \leq 1}} \Delta_i f \Delta_j g.$$

Bony's decomposition gives

$$fg = T_f g + T_g f + R(f, g).$$

Finally, we state Bernstein's inequality. For a proof of this inequality, see [6].

**Lemma 2.1.2.** (*Bernstein's inequality*) Let  $r_1$  and  $r_2$  satisfy  $0 < r_1 < r_2 < \infty$ , and let  $p$  and  $q$  satisfy  $1 \leq p \leq q \leq \infty$ . There exists a positive constant  $C$

such that for every integer  $k$ , if  $u$  belongs to  $L^p(\mathbb{R}^n)$ , and  $\text{supp } \hat{u} \subset B(0, r_1\lambda)$ , then

$$\sup_{|\alpha|=k} \|\partial_\alpha u\|_{L^q} \leq C^k \lambda^{k+n(\frac{1}{p}-\frac{1}{q})} \|u\|_{L^p}. \quad (2.1.1)$$

Furthermore, if  $\text{supp } \hat{u} \subset C(0, r_1\lambda, r_2\lambda)$ , then

$$C^{-k} \lambda^k \|u\|_{L^p} \leq \sup_{|\alpha|=k} \|\partial^\alpha u\|_{L^p} \leq C^k \lambda^k \|u\|_{L^p}. \quad (2.1.2)$$

## 2.2 Function spaces

We now define several useful function spaces.

**Definition 2.2.1.** Assume  $s \in (0, 1)$ . We define the Holder space  $C^s(\mathbb{R}^n)$  to be the space of bounded functions  $f$  on  $\mathbb{R}^n$  such that there exists a constant  $C$  with

$$|f(x) - f(y)| \leq C|x - y|^s \quad (2.2.1)$$

for all  $x$  and  $y$  in  $\mathbb{R}^n$ . For  $s$  a non-integer number greater than 1, we define  $C^s$  to be the set of functions  $f$  such that, for all multi-indices  $\alpha$  with  $|\alpha| \leq [s]$ ,  $\partial^\alpha f$  belongs to  $C^{s-[s]}$ . We define the  $C^s$ -norm as follows:

$$\|f\|_{C^s} = \sum_{|\alpha| \leq [s]} \left( \|\partial^\alpha f\|_{L^\infty} + \sup_{x \neq y} \frac{|\partial^\alpha f(x) - \partial^\alpha f(y)|}{|x - y|^{s-[s]}} \right). \quad (2.2.2)$$

**Definition 2.2.2.** Let  $s \in \mathbb{R}$ . We define the Zygmund space  $C_*^s(\mathbb{R}^n)$  to be the space of tempered distributions  $f$  on  $(\mathbb{R}^n)$  such that

$$\|f\|_{C_*^s} := \sup_{q \geq -1} 2^{qs} \|\Delta_q f\|_{L^\infty} < \infty.$$

It is well known that the norm on  $C_*^s$  is equivalent to the classical  $C^s$ -norm when  $s$  is not an integer and  $s > 0$ . For a proof of this, see [6], Proposition 2.3.1.

**Definition 2.2.3.** Let  $s \in \mathbb{R}$ ,  $p, q \in [1, \infty]$ . We define the inhomogeneous Besov space  $B_{p,q}^s(\mathbb{R}^n)$  to be the space of tempered distributions  $f$  on  $(\mathbb{R}^n)$  such that

$$\|f\|_{B_{p,q}^s} := \left( \sum_{j=-1}^{\infty} 2^{jq_s} \|\Delta_j f\|_{L^p}^q \right)^{\frac{1}{q}} < \infty.$$

When  $q = \infty$ , write

$$\|f\|_{B_{p,\infty}^s} := \sup_{j \geq -1} 2^{js} \|\Delta_j f\|_{L^p}.$$

**Remark 2.2.1.** Note that  $B_{\infty,\infty}^s = C_*^s$ .

**Definition 2.2.4.** Let  $s \in \mathbb{R}$ ,  $p, q \in [1, \infty]$ . We define the inhomogeneous Triebel-Lizorkin space  $F_{p,q}^s(\mathbb{R}^n)$  to be the space of tempered distributions  $f$  on  $\mathbb{R}^n$  such that

$$\|f\|_{F_{p,q}^s} := \left\| \left( \sum_{j=-1}^{\infty} 2^{jq_s} |\Delta_j f(\cdot)|^q \right)^{\frac{1}{q}} \right\|_{L^p} < \infty.$$

When  $q = \infty$ , write

$$\|f\|_{F_{p,\infty}^s} := \left\| \sup_{j \geq -1} 2^{js} |\Delta_j f(\cdot)| \right\|_{L^p}.$$

It is well known that the space  $F_{p,2}^s(\mathbb{R}^n)$  coincides with the classical Sobolev space  $W^{s,p}(\mathbb{R}^n)$  whenever  $1 < p < \infty$  (see, for example, [7]).

**Definition 2.2.5.** The space of log-Lipschitz functions, denoted by  $LL(\mathbb{R}^n)$ , is the space of bounded functions  $f$  on  $\mathbb{R}^n$  such that

$$\|f\|_{LL} := \|f\|_{L^\infty} + \sup_{|x-y| \leq 1} \frac{|f(x) - f(y)|}{|x-y|(1 - \log|x-y|)} < \infty.$$

We now define  $BMO(\mathbb{R}^n)$ , a space first introduced by John and Nirenberg in [10].

**Definition 2.2.6.** Assume  $f$  is a function defined on  $\mathbb{R}^n$ . Then  $f$  has bounded mean oscillation (or, equivalently, belongs to the space  $BMO(\mathbb{R}^n)$ ) if there exists a positive constant  $C$  such that for all balls  $B$  in  $\mathbb{R}^n$ ,

$$\frac{1}{m(B)} \int_B |f(x) - f_B| dx \leq C, \quad (2.2.3)$$

where  $f_B$  denotes the average of  $f$  over  $B$ . The  $BMO$ -seminorm of  $f$  is defined to be the smallest constant  $C$  satisfying (2.2.3).

It is easy to see that all bounded functions on  $(\mathbb{R}^n)$  are also in  $BMO(\mathbb{R}^n)$ , and  $\|f\|_{BMO} \leq 2\|f\|_{L^\infty}$ .

We will also make use of the space  $bmo(\mathbb{R}^n)$  which is a local version of the space  $BMO(\mathbb{R}^n)$ .

**Definition 2.2.7.** The space  $bmo(\mathbb{R}^n)$  is the set of all locally integrable functions  $f$  on  $\mathbb{R}^n$  such that

$$\sup_{|B| \leq 1} \frac{1}{m(B)} \int_B |f(x) - f_B| dx + \sup_{|B|=1} \int_B |f(x)| dx \leq C. \quad (2.2.4)$$

It is clear from the definition that  $L^p(\mathbb{R}^n) \cap BMO(\mathbb{R}^n)$  continuously embeds into  $bmo(\mathbb{R}^n)$ . Moreover, the space  $bmo(\mathbb{R}^n)$  is equivalent to the Triebel-Lizorkin space  $F_{\infty,2}^0(\mathbb{R}^n)$  (see [19]).

We define  $VMO(\mathbb{R}^n)$  and  $vmo(\mathbb{R}^n)$ , first introduced by Sarason in [15].

**Definition 2.2.8.** The space  $VMO(\mathbb{R}^n)$  is the closure of the space  $C_c^\infty(\mathbb{R}^n)$  in the  $BMO$ -norm. Similarly,  $vmo(\mathbb{R}^n)$  is the closure of the space  $C_c^\infty(\mathbb{R}^n)$  in the  $bmo$ -norm.

We now state several important embeddings which we use in what follows. For proofs of these embeddings, we refer the reader to [7] or [19].

**Proposition 2.2.1.** *The following embeddings hold:*

- (i)  $B_{p,q}^s \hookrightarrow F_{p,q}^s$  for  $s \in \mathbb{R}$ ,  $0 < q < p \leq \infty$ .
- (ii)  $F_{p,q}^s \hookrightarrow B_{p,q}^s$  for  $s \in \mathbb{R}$ ,  $0 < p < q \leq \infty$ .
- (iii)  $B_{p,q_1}^s \hookrightarrow B_{p,q_2}^s$  for  $s \in \mathbb{R}$ ,  $q_1 < q_2$ .
- (iv)  $F_{p,q_1}^s \hookrightarrow F_{p,q_2}^s$  for  $s \in \mathbb{R}$ ,  $q_1 < q_2$ .
- (v)  $B_{p,q_1}^s \hookrightarrow B_{p,q_2}^t$  for  $t < s$  and for all  $q_1, q_2$ .
- (vi)  $F_{p,q_1}^s \hookrightarrow F_{p,q_2}^t$  for  $t < s$  and for all  $q_1, q_2$ .

## 2.3 Properties of ideal incompressible fluids

In this section we state some properties of ideal incompressible fluids. We will make use of these properties in what follows. We begin with the

Biot-Savart law.

### 2.3.1 The Biot-Savart law

The Biot-Savart law states that the velocity of an ideal incompressible fluid can be uniquely determined by the vorticity as long as the vorticity has sufficient decay at infinity:

**Theorem 2.3.1.** (*Biot-Savart law*) Assume  $\omega$  belongs to  $L^a(\mathbb{R}^2)$  for some  $a < 2$ . If  $b > \frac{2a}{2-a}$ , then there exists a unique divergence-free velocity vector field  $v$  in  $L^a(\mathbb{R}^2) + L^b(\mathbb{R}^2)$  with vorticity equal to  $\omega$ . If  $E_2$  denotes the fundamental solution of the Laplacian in dimension 2, (that is,  $E_2 = (2\pi)^{-1} \log|x|$ ), then

$$v = (-\partial_2 E_2 * \omega, \partial_1 E_2 * \omega). \quad (2.3.1)$$

*Proof.* Write  $\tilde{v}^i = (-1)^i \partial_j E_2 * \omega(v)$ . We will show that  $\omega(\tilde{v}) = \omega(v)$  and  $\operatorname{div} \tilde{v} = 0$ . This will imply by Lemma 2.3.5 in Section 2.3.3 that  $\tilde{v} = v$ . To show that  $\omega(\tilde{v}) = \omega(v)$ , we write

$$\begin{aligned} \omega(\tilde{v}) &= \partial_1 \partial_1 E_2 * \omega(v) + \partial_2 \partial_2 E_2 * \omega(v) \\ &= \Delta E_2 * \omega(v) = \omega(v). \end{aligned} \quad (2.3.2)$$

Moreover,  $\operatorname{div} \tilde{v} = -\partial_1 \partial_2 E_2 * \omega + \partial_2 \partial_1 E_2 * \omega = 0$ . Therefore,  $v = \tilde{v} + \tilde{p}$ , where the coefficients of  $\tilde{p}$  are harmonic polynomials. Since  $\omega$  belongs to  $L^a(\mathbb{R}^2)$  for  $a < 2$ , and  $|\partial_i E_2(x)| \leq \chi \frac{1}{|x|} + (1 - \chi) \frac{1}{|x|}$  belongs to  $L^1(\mathbb{R}^2) + L^s(\mathbb{R}^2)$  for  $s > 2$ ,  $\tilde{v}$  belongs to  $L^a(\mathbb{R}^2) + L^b(\mathbb{R}^2)$  by Young's inequality. This implies that  $\tilde{p} = 0$  and  $\tilde{v} = v$ . This completes the proof.  $\square$

If we take the gradient of each side of (2.3.1), we get

$$\nabla v = (-\nabla \partial_2 E_2 * \omega, \nabla \partial_1 E_2 * \omega), \quad (2.3.3)$$

and we see that  $\nabla v$  can be written as a Calderon-Zygmund operator acting on  $\omega$ . We can therefore use mapping properties of Calderon-Zygmund operators to determine information about  $\nabla v$  from our assumptions on  $\omega$ . For example, it is well known that Calderon-Zygmund operators map  $L^p(\mathbb{R}^n)$  to  $L^p(\mathbb{R}^n)$  for  $1 < p < \infty$ . These operators are not bounded from  $L^\infty(\mathbb{R}^n)$  to  $L^\infty(\mathbb{R}^n)$ ; however, they continuously map  $L^\infty(\mathbb{R}^n)$  into  $BMO(\mathbb{R}^n)$  (see [18] or [16] for details).

### 2.3.2 The vorticity of an ideal incompressible fluid

The goal of this section is to introduce properties of weak solutions to the two-dimensional Euler equations. We use these properties in Chapter 3.

We begin this section by taking the curl of (E) and using the definition of vorticity in  $\mathbb{R}^2$  given by  $\omega = \partial_1 v^2 - \partial_2 v^1$  to get the two-dimensional vorticity equation:

$$(V) \quad \begin{cases} \partial_t \omega + v \cdot \nabla \omega = 0, \\ \omega|_{t=0} = \omega^0. \end{cases}$$

When  $v$  is smooth, we can easily conclude from (V) that the  $L^p$ -norm of vorticity is conserved in two dimensions for all  $p \in [1, \infty)$ . To see this, we multiply (V) by  $|\omega|^{r-2} \omega$  and observe that

$$\partial_t |\omega|^r = r |\omega|^{r-2} \omega (\partial_t \omega) = -r |\omega|^{r-2} \omega (v \cdot \nabla \omega). \quad (2.3.4)$$

Integrating both sides of (2.3.4), we get

$$\begin{aligned} \partial_t \int_{\mathbb{R}^2} |\omega|^r dx &= - \int_{\mathbb{R}^2} u \cdot r |\omega|^{r-2} \omega \nabla \omega \\ &= \int_{\mathbb{R}^2} (\operatorname{div} u) |\omega|^r dx = 0. \end{aligned} \tag{2.3.5}$$

We can draw the same conclusions using the flow  $g : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ . The flow is defined by the following ordinary differential equation:

$$\begin{aligned} \partial_t g(t, x) &= v(t, g(t, x)), \\ g(0, x) &= x. \end{aligned} \tag{2.3.6}$$

When  $v$  is smooth, one can infer from the divergence-free condition on  $v$  that the flow is measure-preserving (see, for example, chapter 1 of [2]). Moreover, we can rewrite (V) using (2.3.6) and the chain rule as

$$\partial_t (\omega(t, g(t, x))) = 0, \tag{2.3.7}$$

which implies that the vorticity is conserved along flow lines and once again that the  $L^p$ -norms of  $\omega$  are conserved over time.

In this section we discuss why these properties hold in a weaker setting. Throughout the discussion, we assume only that  $\omega^0$  belongs to the space  $L^1(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ . We closely follow chapter 8 of [2], and we refer the reader to [2] for further details. We will outline the proof of the following theorem:

**Theorem 2.3.2.** *Given  $\omega^0 \in L^1(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ , there exists a unique weak solution to (V) in the sense that there exists a unique pair  $(v, \omega)$  satisfying:*

- (i)  $\omega$  belongs to  $L^\infty([0, T]; L^1(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2))$ ,
- (ii)  $\omega = \omega(v)$  and  $v = K * \omega$ , where  $K$  is the Biot-Savart kernel,

(iii) for all  $\varphi \in C^1([0, T]; C_0^1(\mathbb{R}^2))$ ,

$$\int_{\mathbb{R}^2} \varphi(T, x) \omega(T, x) dx - \int_{\mathbb{R}^2} \varphi(0, x) \omega(0, x) dx = \int_0^T \int_{\mathbb{R}^2} (\partial_t \varphi + v \cdot \nabla \varphi) \omega dx dt. \quad (2.3.8)$$

In fact, uniqueness was proved by Yudovich in [23]. In what follows, we outline a proof for existence of such solutions. From this existence proof, we are able to show that several of the properties satisfied by classical solutions carry over to the weak solution  $(v, \omega)$ .

*Proof.* We begin by mollifying the initial vorticity. We consider a sequence of solutions  $(\omega_n)$  to (V) with initial data  $S_n \omega^0$ , and with velocity  $v_n$  determined uniquely by the Biot-Savart law. We infer from the discussion above that

$$\omega_n(t) = S_n \omega^0(g_n(t)^{-1}(x)), \quad (2.3.9)$$

where  $g_n(t)$  satisfies

$$\begin{aligned} \partial_t g_n(t, x) &= v_n(t, g_n(t, x)), \\ g_n(0, x) &= x. \end{aligned} \quad (2.3.10)$$

Moreover, by Lemma 2.3.9 in Section 2.3.3 and (2.3.10), we have

$$|g_n(t)^{-1}(x) - x| = \left| \int_0^t v_n(\tau, g_n(\tau, x)) d\tau \right| \leq CT. \quad (2.3.11)$$

This uniform bound, combined with Holder estimates on  $g_n^{-1}$  (see Lemma 8.2 of [2]), imply that the sequence  $(g_n^{-1})$  is equicontinuous on  $K \times [0, T]$  for  $K \subseteq \mathbb{R}^2$  compact. We apply the Arzela-Ascoli theorem and conclude that

there exists a subsequence converging uniformly on  $K \times [0, T]$  to a function which we denote by  $g^{-1}$ .

We now claim that  $\omega_n^0(g_n(t)^{-1})$  converges to  $\omega^0(g(t)^{-1})$  in  $L^1(\mathbb{R}^2)$ , and that  $\omega^0(g(t)^{-1})$  solves the weak formulation of (V). To see this, we first need to establish that  $g(t)^{-1}$  is indeed a measure-preserving map for each  $t$  in  $[0, T]$ .

**Lemma 2.3.3.** *For every  $t$  in  $[0, T]$ , the map  $g(t)^{-1}$  established above is measure-preserving on  $\mathbb{R}^2$ .*

*Proof.* We follow the approach in [2] and prove that  $\int_{\mathbb{R}^2} f(g(t)^{-1}(x))dx = \int_{\mathbb{R}^2} f(x)dx$  for all  $f$  in  $L^1(\mathbb{R}^2)$ . We assume that  $f$  is continuous with compact support and use density of  $C_c(\mathbb{R}^2)$  in  $L^1(\mathbb{R}^2)$ . Because  $f$  is in  $C_c(\mathbb{R}^2)$ , and  $g_n(t)^{-1}$  converges uniformly to  $g(t)^{-1}$  on compact sets, we have that  $f(g_n(t)^{-1})$  converges to  $f(g(t)^{-1})$  pointwise. We apply Lebesgue's dominated convergence theorem and conclude that

$$\int_{\mathbb{R}^2} f(g(t)^{-1})dx = \lim_{n \rightarrow \infty} \int_{\mathbb{R}^2} f(g_n(t)^{-1})dx = \int_{\mathbb{R}^2} f dx. \quad (2.3.12)$$

We use the Riesz representation theorem and (2.3.12) to conclude that  $|B|$  and  $|g(t)^{-1}(B)|$  are equivalent on Borel sets. Therefore,  $g(t)^{-1}$  is measure-preserving.  $\square$

We now use Lemma 2.3.3 to show that  $\omega_n^0(g_n(t)^{-1})$  converges to  $\omega^0(g(t)^{-1})$  in  $L^1(\mathbb{R}^2)$ . We write

$$\begin{aligned} \|\omega_n^0(g_n(t)^{-1}) - \omega^0(g(t)^{-1})\|_{L^1} &\leq \|\omega_n^0(g_n(t)^{-1}) - \omega^0(g_n(t)^{-1})\|_{L^1} \\ &+ \|\omega^0(g_n(t)^{-1}) - \omega^0(g(t)^{-1})\|_{L^1}. \end{aligned} \quad (2.3.13)$$

The first term on the right hand side of (2.3.13) is equal to  $\|\omega_n^0 - \omega^0\|_{L^1}$ , which converges to 0 as  $n$  approaches infinity by our construction of the smooth initial data. We break the second term on the right hand side of (2.3.13) down as in [2]; we let  $\omega_m^0$  be a continuous function with compact support on  $\mathbb{R}^2$  such that  $\|\omega^0 - \omega_m^0\|_{L^1} < \frac{1}{m}$ , and we write

$$\begin{aligned} \|\omega^0(g_n(t)^{-1}) - \omega^0(g(t)^{-1})\|_{L^1} &\leq \|\omega^0(g_n(t)^{-1}) - \omega_m^0(g_n(t)^{-1})\|_{L^1} \\ &\quad + \|\omega_m^0(g_n(t)^{-1}) - \omega_m^0(g(t)^{-1})\|_{L^1} + \|\omega_m^0(g(t)^{-1}) - \omega^0(g(t)^{-1})\|_{L^1}. \end{aligned} \tag{2.3.14}$$

Since  $g_n(t)^{-1}$  and  $g(t)^{-1}$  are measure-preserving, we can choose  $m$  sufficiently large to make the first and third terms on the right hand side of (2.3.14) small. For the middle term, for fixed, sufficiently large  $m$ , we use sequential continuity of  $\omega_m^0$  to apply Lebesgue's dominated convergence theorem. This completes the argument that  $\omega_n^0(g_n(t)^{-1})$  converges to  $\omega^0(g(t)^{-1})$  in  $L^1(\mathbb{R}^2)$ .

Using this convergence, we are also able to show that  $v_n(t)$  converges to  $v(t)$  uniformly on compact sets for fixed  $t \in [0, T]$ , where  $v_n(t)$  and  $v(t)$  are determined uniquely from  $\omega_n$  and  $\omega$  using the Biot-Savart law. The strategy of the proof is to write the difference of the two velocities as a convolution of the Biot-Savart kernel with the difference of the two vorticities, to apply Young's inequality to the convolution, and to utilize convergence of the sequence  $(\omega_n)$  to  $\omega$  in  $L^1(\mathbb{R}^2)$ . We refer the reader to [2] for a detailed proof.

Finally, we use convergence of  $(\omega_n)$  to  $\omega$  in  $L^1(\mathbb{R}^2)$  and uniform convergence of  $v_n(t)$  to  $v(t)$  on compact sets to show that  $\omega(t) = \omega^0(g(t)^{-1})$  solves our weak formulation. We recall the definition of a weak solution:

**Definition 2.3.1.** Given  $\omega^0 \in L^1(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ , the pair  $(v, \omega)$  is a weak solution to (V) if the following three conditions hold:

- (i)  $\omega$  belongs to  $L^\infty([0, T]; L^1(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2))$ ,
- (ii)  $\omega = \omega(v)$  and  $v = K * \omega$ , where  $K$  is the Biot-Savart kernel,
- (iii) for all  $\varphi \in C^1([0, T]; C_0^1(\mathbb{R}^2))$ ,

$$\int_{\mathbb{R}^2} \varphi(T, x) \omega(T, x) dx - \int_{\mathbb{R}^2} \varphi(0, x) \omega(0, x) dx = \int_0^T \int_{\mathbb{R}^2} (\partial_t \varphi + v \cdot \nabla \varphi) \omega dx dt. \quad (2.3.15)$$

A straightforward computation (see Proposition 8.1 of [2]) shows that every smooth solution of (V) solves the weak formulation. We must show that  $\omega^0(g(t)^{-1})$  satisfies (iii) as well. We will show that

$$\begin{aligned} \int_{\mathbb{R}^2} \varphi(T, x) \omega_n(T, x) dx - \int_{\mathbb{R}^2} \varphi(0, x) \omega_n(0, x) dx \\ = \int_0^T \int_{\mathbb{R}^2} (\partial_t \varphi + v_n \cdot \nabla \varphi) \omega_n dx dt \end{aligned} \quad (2.3.16)$$

converges to

$$\int_{\mathbb{R}^2} \varphi(T, x) \omega(T, x) dx - \int_{\mathbb{R}^2} \varphi(0, x) \omega(0, x) dx = \int_0^T \int_{\mathbb{R}^2} (\partial_t \varphi + v \cdot \nabla \varphi) \omega dx dt \quad (2.3.17)$$

as  $n$  approaches infinity. The term  $\int_{\mathbb{R}^2} \varphi(T, x) \omega_n(T, x) dx$  clearly converges to  $\int_{\mathbb{R}^2} \varphi(T, x) \omega(T, x) dx$  because  $\varphi$  is in  $C_0^1(\mathbb{R}^2)$  and  $\omega_n$  converges to  $\omega$  in  $L^1(\mathbb{R}^2)$ ; similarly  $\int_{\mathbb{R}^2} \varphi(0, x) \omega_n(0, x) dx$  converges to  $\int_{\mathbb{R}^2} \varphi(0, x) \omega(0, x) dx$ , and  $\int_0^T \int_{\mathbb{R}^2} \partial_t \varphi \omega_n dx dt$  converges to  $\int_0^T \int_{\mathbb{R}^2} \partial_t \varphi \omega dx dt$ . To handle the nonlinearity, we observe that  $\omega_n$  is uniformly bounded independent of  $n$  by  $C \|\omega^0\|_{L^\infty}$ , and

$\|v_n\|_{L^\infty} \leq \|v\|_{L^\infty} + \epsilon$  for large  $n$ . Furthermore, there exists a subsequence of  $(v_n \cdot \nabla \varphi) \omega_n$  converging pointwise to  $(v \cdot \nabla \varphi) \omega$ . We apply Lebesgue's dominated convergence theorem, and we conclude that  $\omega$  is a weak solution to (V).  $\square$

**Remark 2.3.1.** From this discussion we draw several conclusions. Most importantly, we are able to conclude that for a solution  $\omega$  to the weak formulation given by Definition 2.3.1, the solution possesses a flow  $g$  such that  $g(t)$  and  $g(t)^{-1}$  are measure-preserving on  $\mathbb{R}^2$ . Moreover, the weak solution  $\omega$  satisfies  $\omega(t) = \omega^0(g(t)^{-1})$  by construction. In fact, Yudovich shows in [23] that these properties hold for our solution if we assume only that  $\omega^0$  belongs to  $L^p(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$  for some  $p \in [1, \infty)$ . One can prove this using an approximation argument similar to that given above. We refer the reader to [23] for details.

### 2.3.3 A few technical lemmas

In this section we state and prove a few technical lemmas. We frequently make use of these lemmas in what follows.

We begin with Osgood's Lemma. We refer the reader to [5], pg. 92, for a detailed proof.

**Lemma 2.3.4.** (*Osgood's Lemma*) *Let  $\rho$  be a measurable positive function,  $\gamma$  a positive locally integrable function, and  $\mu$  a continuous, increasing function. Assume that  $a$  is some positive real number and that*

$$\rho(t) \leq a + \int_{t_0}^t \gamma(s) \mu(\rho(s)) ds. \quad (2.3.18)$$

If we define  $M(x) = \int_x^1 \frac{dr}{\mu(r)}$ , then we have

$$-M(\rho(t)) + M(a) \leq \int_{t_0}^t \gamma(s) ds. \quad (2.3.19)$$

**Lemma 2.3.5.** *Two vector fields whose coefficients are tempered distributions and with the same vorticity and divergence are equal up to a vector field with harmonic polynomials as coefficients.*

*Proof.* We have

$$\partial_i v^j = \partial_j v^i + \omega(v), \quad (2.3.20)$$

with  $i \neq j$ . Taking the partial derivative with respect to the  $i$ th variable of both sides and summing over  $i = 1, 2$  gives

$$\Delta v^i = \partial_j \operatorname{div} v + \sum_i \partial_i \omega(v). \quad (2.3.21)$$

Therefore, if  $\operatorname{div} v = \operatorname{div} \tilde{v}$  and  $\omega(v) = \omega(\tilde{v})$ , then  $\Delta(v - \tilde{v}) = 0$ , and  $v - \tilde{v}$  is a harmonic polynomial. This completes the proof.  $\square$

**Lemma 2.3.6.** *If  $v$  is a divergence-free vector field in  $L^2(\mathbb{R}^2)$  with vorticity  $\omega$  in  $L^2(\mathbb{R}^2)$  then  $v$  belongs to  $H^1(\mathbb{R}^2)$  and  $\|\nabla v\|_{L^2} = \|\omega\|_{L^2}$ .*

*Proof.* Because  $v$  belongs to  $L^2(\mathbb{R}^2)$ , the Fourier transform of  $v$  belongs to  $L^2(\mathbb{R}^2)$ , and  $\widehat{\partial_j v^i}(\xi) = \xi_j \hat{v}^i(\xi)$  is defined pointwise almost everywhere. There-

fore, we can write

$$\begin{aligned}
|\widehat{\nabla}v(\xi)|^2 &= \xi_1^2(\hat{v}^1(\xi))^2 + \xi_1^2(\hat{v}^1(\xi))^2 + \xi_2^2(\hat{v}^1(\xi))^2 + \xi_2^2(\hat{v}^2(\xi))^2 \\
&= \xi_1^2(\hat{v}^1(\xi))^2 + \xi_2^2(\hat{v}^2(\xi))^2 + 2\xi_1\xi_2\hat{v}^1(\xi)\hat{v}^2(\xi) \\
&+ \xi_1^2(\hat{v}^2(\xi))^2 + \xi_2^2(\hat{v}^1(\xi))^2 - 2\xi_1\xi_2\hat{v}^1(\xi)\hat{v}^2(\xi) \\
&= (\xi_1\hat{v}^1(\xi) + \xi_2\hat{v}^2(\xi))^2 + (\xi_1\hat{v}^2(\xi) - \xi_2\hat{v}^1(\xi))^2 \\
&= \left(\widehat{\operatorname{div}v}(\xi)\right)^2 + (\hat{\omega}(\xi))^2 = (\hat{\omega}(\xi))^2
\end{aligned} \tag{2.3.22}$$

for almost every  $\xi \in \mathbb{R}^2$ . Thus  $\|\nabla v\|_{L^2} = \|\widehat{\nabla}v\|_{L^2} = \|\hat{\omega}\|_{L^2} = \|\omega\|_{L^2}$ . This completes the proof.  $\square$

In Section 2.3.1, we showed that whenever  $\omega$  belongs to  $L^p(\mathbb{R}^2)$  for  $p < 2$ , the velocity  $v$  can be uniquely determined from  $\omega$  using the Biot-Savart law. If we only assume  $\omega$  belongs to  $L^p(\mathbb{R}^2)$  for some  $p \geq 2$ , we cannot *determine*  $v$  from  $\omega$ ; however, if  $v$  is a divergence-free vector field in  $L^p(\mathbb{R}^2)$  for some  $p < \infty$  with vorticity  $\omega \in L^q(\mathbb{R}^2)$  for  $q \in [2, \infty)$ , we can still write  $v$  as the Biot-Savart kernel convolved with  $\omega$ . We have the following lemma.

**Lemma 2.3.7.** *Let  $p \in (1, \infty)$  and  $q \in [2, \infty)$ , and assume  $v$  is a divergence-free vector field in  $L^p(\mathbb{R}^2)$  with  $\omega = \omega(v)$  in  $L^q(\mathbb{R}^2)$ . Then  $v = (-\partial_2 E_2 * \omega, \partial_1 E_2 * \omega)$ .*

*Proof.* Let  $\psi$  be a tempered distribution on  $\mathbb{R}^2$  satisfying  $v = (-\partial_2\psi, \partial_1\psi)$ , so that  $\Delta\psi = \omega$ . (To see that such a tempered distribution exists, we refer the reader to Corollary 1.2.2 on page 7 of [5].) Let  $\psi' = \Delta^{-1}\omega$ . Then, if

$v' = (-\partial_2\psi', \partial_1\psi')$ , we have  $\operatorname{div} v' = 0$  and  $\omega(v') = \Delta\psi' = \omega$ . Therefore,  $v$  and  $v'$  have the same divergence and vorticity. By Lemma 2.3.5, we can write

$$v - v' = q, \quad (2.3.23)$$

where  $q$  is a vector field with polynomials as coefficients. Using the equalities  $v' = \nabla^\perp\psi = \nabla^\perp\Delta^{-1}\omega$ , we see that for  $r \in [\max\{p, q\}, \infty)$ ,

$$\begin{aligned} \|v'\|_{L^r} &= \|\Delta_{-1}v'\|_{L^r} + \sum_{j \geq 0} \|\Delta_j v'\|_{L^r} \\ &\leq C\|\Delta_{-1}v'\|_{L^p} + C \sum_{j \geq 0} 2^{2j(\frac{1}{q} - \frac{1}{r})-j} \|\Delta_j \nabla v'\|_{L^q} \\ &= C\|\Delta_{-1}\nabla^\perp\Delta^{-1}\omega\|_{L^p} + C \sum_{j \geq 0} 2^{2j(\frac{1}{q} - \frac{1}{r})-j} \|\Delta_j \nabla \nabla^\perp \Delta^{-1}\omega\|_{L^q} \quad (2.3.24) \\ &\leq C\|\Delta_{-1}\nabla^\perp\Delta^{-1}\nabla v\|_{L^p} + C\|\omega\|_{L^q} \\ &\leq C\|v\|_{L^p} + C\|\omega\|_{L^q}, \end{aligned}$$

where we repeatedly used Bernstein's inequality, and we used boundedness of Calderon-Zygmund operators on  $L^s(\mathbb{R}^2)$  for  $s \in (1, \infty)$  to get the second-to-last and last inequalities. Since  $v \in L^p(\mathbb{R}^2)$  and  $v' \in L^r(\mathbb{R}^2)$ , it follows that  $q = 0$  in (2.3.23). Therefore,  $v = v' = \nabla^\perp\Delta^{-1}\omega$ . This completes the proof.  $\square$

**Lemma 2.3.8.** *Let  $v$  be a divergence-free vector field in  $L^2(\mathbb{R}^2)$  with vorticity  $\omega$  in  $L^2(\mathbb{R}^2)$ . There exists an absolute constant  $C$  such that for all  $q \geq 0$  (that is, avoiding the low frequencies),*

$$\|\Delta_q \nabla v\|_{L^\infty} \leq C \|\Delta_q \omega\|_{L^\infty}.$$

*Proof.* Since  $v$  is a divergence-free vector field in  $L^2(\mathbb{R}^2)$ , and  $\omega \in L^2(\mathbb{R}^2)$ , it

follows by Lemma 2.3.7 that  $\nabla \Delta_q v = \nabla \nabla^\perp \Delta^{-1} \Delta_q \omega$ . We can then write

$$\|\Delta_q \nabla v\|_{L^\infty} \leq C \sup_{i,j} \|\Delta_q \partial_i \partial_j \Delta^{-1} \omega\|_{L^\infty}.$$

But,

$$\begin{aligned} \|\Delta_q \partial_i \partial_j \Delta^{-1} \omega\|_{L^\infty} &= \|\mathcal{F}^{-1}(\varphi_q(\xi) \frac{\xi_i \xi_j}{|\xi|^2} \widehat{\omega}(\xi))\|_{L^\infty} \\ &= \|\mathcal{F}^{-1}(\varphi_q(\xi) h_q(\xi) \widehat{\omega}(\xi))\|_{L^\infty} = \|\Delta_q(\check{h}_q(\xi) * \omega)\|_{L^\infty}, \end{aligned}$$

where

$$h_q(\xi) = \chi(2^{-3-q}\xi)(1 - \chi(2^{-q+1}\xi)) \frac{\xi_i \xi_j}{|\xi|^2},$$

$\chi$  and  $\varphi$  being defined in Lemma 2.1.1. Observe that  $h_q = 1$  on the support of  $\varphi_q$ . Because  $h_q(\xi) = h(2^{-q}\xi)$ , where

$$h(\xi) = \chi(2^{-3}\xi)(1 - \chi(2\xi)) \frac{\xi_i \xi_j}{|\xi|^2},$$

$\check{h}_q(x) = 2^{2q} \check{h}(2^q x)$ , and thus by a change of variables,  $\|\check{h}_q\|_{L^1} = \|\check{h}\|_{L^1} = C$ .

Then using Young's inequality,

$$\begin{aligned} \|\Delta_q(\check{h}_q(\xi) * \omega)\|_{L^\infty} &= \|\check{h}_q(\xi) * \Delta_q \omega\|_{L^\infty} \leq \|\check{h}_q\|_{L^1} \|\Delta_q \omega\|_{L^\infty} \\ &\leq C \|\Delta_q \omega\|_{L^\infty}, \end{aligned}$$

which completes the proof.  $\square$

**Lemma 2.3.9.** *Assume  $\omega$  is a  $C^\infty$  solution to (V) with  $\omega^0 \in L^p(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$  for some  $p \in [1, 2)$ , and with  $v$  determined from  $\omega$  using the Biot-Savart law. Then  $v$  belongs to  $L_{loc}^\infty(\mathbb{R}^+; L^\infty(\mathbb{R}^2))$ , and  $\|v(t)\|_{L^\infty} \leq C \|\omega^0\|_{L^p \cap L^\infty}$ .*

*Proof.* The proof of Lemma 2.3.9 closely follows arguments in [21], Theorem 3.1. We break  $v$  into low and high frequencies and write

$$\begin{aligned}
\|v(t)\|_{L^\infty} &\leq \|\Delta_{-1}v(t)\|_{L^\infty} + \sum_{j \geq 0} \|\Delta_j v(t)\|_{L^\infty} \\
&\leq \|\Delta_{-1}v(t)\|_{L^\infty} + \sum_{j \geq 0} 2^{-j} \|\Delta_j \nabla v(t)\|_{L^\infty} \\
&\leq \|\Delta_{-1}v(t)\|_{L^\infty} + C\|\omega^0\|_{L^\infty},
\end{aligned} \tag{2.3.25}$$

where we used Lemma 2.3.8, Bernstein's inequality, and conservation of vorticity along flow lines for the high frequency terms. To bound the low frequency term, we use the Biot-Savart law. Let  $\chi$  be a smooth bump function with support in the unit ball. Write

$$\begin{aligned}
\|\Delta_{-1}v(t)\|_{L^\infty} &\leq \|\chi \partial_i E_2 \Delta_{-1} \omega(t)\|_{L^\infty} + \|(1 - \chi) \partial_i E_2 \Delta_{-1} \omega(t)\|_{L^\infty} \\
&\leq \|\chi \partial_i E_2\|_{L^1} \|\Delta_{-1} \omega(t)\|_{L^\infty} + \|(1 - \chi) \partial_i E_2\|_{L^{q_0}} \|\Delta_{-1} \omega(t)\|_{L^{p_0}} \\
&\leq C \|\Delta_{-1} \omega(t)\|_{L^{p_0}} \leq C \|\omega^0\|_{L^{p_0}},
\end{aligned} \tag{2.3.26}$$

where  $\frac{1}{p_0} + \frac{1}{q_0} = 1$ . Again, we used conservation of the vorticity along flow lines. This completes the proof.  $\square$

The following lemma and proof can be found in [21], Theorem 3.1.

**Lemma 2.3.10.** *Assume  $v$  is a solution to (E) and  $\omega(v^0) = \omega^0$  belongs to  $L^p(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$  for some  $p < \infty$ . Then  $v$  belongs to  $L^1_{loc}(\mathbb{R}^+; LL(\mathbb{R}^2))$ .*

*Proof.* We begin by showing  $v$  is bounded on  $\mathbb{R}^2$ . Write

$$\begin{aligned}
\|v(t)\|_{L^\infty} &\leq \|\Delta_{-1}v(t)\|_{L^\infty} + \sum_{j \geq 0} \|\Delta_j v(t)\|_{L^\infty} \\
&\leq \|\Delta_{-1}v(t)\|_{L^\infty} + \sum_{j \geq 0} 2^{-j} \|\Delta_j \nabla v(t)\|_{L^\infty} \\
&\leq \|\Delta_{-1}v(t)\|_{L^\infty} + C\|\omega(t)\|_{L^\infty} \\
&\leq C\|\Delta_{-1}v(t)\|_{L^p} + C\|\omega^0\|_{L^\infty},
\end{aligned}$$

where we used Bernstein's inequality, Lemma 2.3.8, and conservation of the  $L^\infty$ -norm of  $\omega$  over time by Remark 2.3.1. To bound  $\|\Delta_{-1}v(t)\|_{L^p}$ , we use the proof of Lemma 6.2 in [20]. We first bound  $\|\Delta_{-1}v(t)\|_{L^p}$  by  $\|v(t)\|_{L^p}$ . Then, because the Weyl projection operator is bounded from  $L^p(\mathbb{R}^2)$  to  $L^p(\mathbb{R}^2)$ , we can write

$$\begin{aligned}
\|v(t)\|_{L^p} - \|v^0\|_{L^p} &\leq \int_0^t \|-v \cdot \nabla v(\tau) - \nabla p(\tau)\|_{L^p} d\tau \\
&\leq C \int_0^t \|v \cdot \nabla v(\tau)\|_{L^p} d\tau \\
&\leq C \int_0^t \|v(\tau)\|_{L^\infty} \|\nabla v(\tau)\|_{L^p} d\tau \\
&\leq C\|\omega^0\|_{L^p} \int_0^t \|v(\tau)\|_{L^\infty} d\tau \\
&\leq C\|\omega^0\|_{L^p} \int_0^t (\|\Delta_{-1}v(t)\|_{L^\infty} + C\|\omega^0\|_{L^p}) d\tau \\
&\leq Ct\|\omega^0\|_{L^p} + C_1\|\omega^0\|_{L^p} \int_0^t \|v(t)\|_{L^p} d\tau,
\end{aligned} \tag{2.3.27}$$

where we used the boundedness of Calderon-Zygmund operators on  $L^p(\mathbb{R}^2)$ , as well as (2.3.30), and conservation of the  $L^p$ -norm of the vorticity. An application of Gronwall's inequality gives

$$\|v(t)\|_{L^p} \leq Ce^{C_1 t \|\omega^0\|_{L^p}}. \tag{2.3.28}$$

We conclude that  $v$  belongs to  $L_{loc}^\infty(\mathbb{R}^+; L^\infty(\mathbb{R}^2))$ . To complete the proof, we write

$$\begin{aligned}
|v(t, x) - v(t, y)| &\leq \sum_{j=-1}^{\infty} |\Delta_j v(t, x) - \Delta_j v(t, y)| \\
&\leq \|\Delta_{-1} \nabla v(t)\|_{L^\infty} |x - y| + \sum_{j=0}^{N-1} \|\Delta_j \nabla v(t)\|_{L^\infty} |x - y| + 2 \sum_{j=N}^{\infty} \|\Delta_j v(t)\|_{L^\infty} \\
&\leq C \|\Delta_{-1} \nabla v(t)\|_{L^p} |x - y| + CN \|\Delta_j \omega(t)\|_{L^\infty} |x - y| + 2 \sum_{j=N}^{\infty} 2^{-j} \|\Delta_j \nabla v(t)\|_{L^\infty} \\
&\leq C \|\omega(t)\|_{L^p} |x - y| + CN \|\omega(t)\|_{L^\infty} |x - y| + C 2^{-N} \|\omega(t)\|_{L^\infty}
\end{aligned} \tag{2.3.29}$$

where we used Bernstein's inequality on the first and third terms, and we used Lemma 2.3.8 on the second and third terms. We also used boundedness of Calderon-Zygmund operators from  $L^p(\mathbb{R}^2)$  to  $L^p(\mathbb{R}^2)$  on the first term. Letting  $N = -\log_2 |x - y|$ , we get

$$\begin{aligned}
|v(t, x) - v(t, y)| &\leq C(\|\omega(t)\|_{L^p} + N \|\omega(t)\|_{L^\infty}) |x - y| + C 2^{-N} \|\omega(t)\|_{L^\infty} \\
&\leq C \|\omega(t)\|_{L^p} (1 - \log_2 |x - y|) |x - y| + C |x - y| \|\omega(t)\|_{L^\infty} \\
&\leq C \|\omega(t)\|_{L^p \cap L^\infty} (1 - \log_2 |x - y|) |x - y|.
\end{aligned}$$

Since the  $L^p$ -norm and  $L^\infty$ -norm of vorticity are conserved over time,  $\|\omega(t)\|_{L^p \cap L^\infty} = \|\omega^0\|_{L^p \cap L^\infty}$ , and the lemma is proved.  $\square$

**Lemma 2.3.11.** *Assume  $v$  is a solution to (E) with  $\omega^0 \in L^p(\mathbb{R}^2)$  for some  $p \in (2, \infty)$ . Then  $v$  belongs to  $L_{loc}^\infty(\mathbb{R}^+; L^\infty(\mathbb{R}^2))$ , and there exist two positive constants  $C$  and  $C_1$  such that  $\|v(t)\|_{L^\infty} \leq C \|\omega^0\|_{L^p} e^{C_1 t \|\omega^0\|_{L^p}}$ .*

*Proof.* The proof of Lemma 2.3.11 is almost identical to the proof that  $v$  belongs to  $L_{loc}^\infty(\mathbb{R}^+; L^\infty(\mathbb{R}^2))$  in Lemma 2.3.10. Here we are assuming  $\omega^0$  belongs to  $L^p(\mathbb{R}^2)$  for fixed  $p \in (2, \infty)$ , but  $\omega^0$  is not necessarily bounded. We therefore treat low frequencies of  $\|v(t)\|_{L^\infty}$  as we did in the proof of Lemma 2.3.11. For high frequencies we modify the proof slightly. We write

$$\begin{aligned}
\|v(t)\|_{L^\infty} &\leq \|\Delta_{-1}v(t)\|_{L^\infty} + \sum_{j \geq 0} \|\Delta_j v(t)\|_{L^\infty} \\
&\leq \|\Delta_{-1}v(t)\|_{L^p} + \sum_{j \geq 0} 2^{j(\frac{2}{p}-1)} \|\Delta_j \nabla v(t)\|_{L^p} \\
&\leq C e^{C_1 t \|\omega^0\|_{L^p}} + C \|\omega^0\|_{L^p},
\end{aligned} \tag{2.3.30}$$

for  $p \in (2, \infty)$ . This completes the proof.  $\square$

## Chapter 3

# An initial value problem for the Euler equations in the plane

### 3.1 Introduction and history

In this chapter, we study regularity of solutions to the Euler equations in the plane with initial velocity in the critical or subcritical Sobolev spaces  $W^{s+1,p}(\mathbb{R}^2)$ , where  $sp = 2$  or  $sp < 2$ , respectively. For the supercritical case, when  $sp > 2$ , Kato and Ponce prove persistence of regularity in [11].

For the critical case, much less is known. In [21] Vishik proves global well-posedness of the two-dimensional Euler equations in the critical Besov space  $B_{p,1}^{s+1}(\mathbb{R}^2)$ . To prove local existence, he observes that  $B_{p,1}^{s+1}(\mathbb{R}^2)$  continuously imbeds into  $B_{\infty,1}^1(\mathbb{R}^2)$ . This implies that  $\nabla v$  belongs to  $B_{\infty,1}^0(\mathbb{R}^2)$  and therefore to  $L^\infty(\mathbb{R}^2)$ . He then uses paradifferential calculus to prove an estimate on the  $B_{p,1}^s$ -norm of the vorticity:

$$\|\omega(t)\|_{B_{p,1}^s} \leq \|\omega^0\|_{B_{p,1}^s} \exp\left(C \int_0^t \|\nabla v(\tau)\|_{L^\infty} d\tau\right). \quad (3.1.1)$$

To give global existence, he proves the following logarithmic estimate on the  $L^\infty$ -norm of  $\nabla v$ :

$$\|\nabla v(t)\|_{L^\infty} \leq C(1 + \log(\|g(t)\|_{lip}\|g^{-1}(t)\|_{lip}))\|\omega^0\|_{B_{\infty,1}^0}. \quad (3.1.2)$$

In [4], Chae proves global well-posedness for the two-dimensional Euler equations in the critical Triebel-Lizorkin spaces  $F_{1,q}^3(\mathbb{R}^2)$ , for  $q \in [1, \infty]$ . Like Vishik, he uses paradifferential calculus to prove the estimate

$$\|\omega(t)\|_{F_{1,q}^2} \leq \|\omega^0\|_{F_{1,q}^2} \exp\left(C \int_0^t \|\nabla v(\tau)\|_{L^\infty} d\tau\right). \quad (3.1.3)$$

In order to obtain a logarithmic estimate like that in (3.1.2), Chae utilizes the following embedding theorem (see, for example, [9], Theorem 2.1):

**Theorem 3.1.1.** *Let  $s_0 > s_1$ ,  $p_0, p_1 \in (0, \infty)$ , and  $q, r \in (0, \infty]$ . If  $s_0 - \frac{n}{p_0} = s_1 - \frac{n}{p_1}$ , then*

$$F_{p_0,q}^{s_0}(\mathbb{R}^n) \hookrightarrow B_{p_1,p_0}^{s_1}(\mathbb{R}^n) \hookrightarrow B_{\infty,p_0}^0(\mathbb{R}^n). \quad (3.1.4)$$

In fact, the first embedding is the interesting part of the theorem. The second embedding follows from a straightforward application of Bernstein's inequality. It follows from Theorem 3.1.1 that  $F_{1,q}^3(\mathbb{R}^2)$  embeds into  $B_{\infty,1}^0(\mathbb{R}^2)$ .

Chae applies the embedding  $B_{\infty,1}^0(\mathbb{R}^2) \hookrightarrow L^\infty(\mathbb{R}^2)$  and boundedness of Calderon-Zygmund operators from  $B_{\infty,1}^0(\mathbb{R}^2)$  into  $B_{\infty,1}^0(\mathbb{R}^2)$  to (3.1.3) and concludes that

$$\|\nabla v(\tau)\|_{L^\infty} \leq \|\nabla v(\tau)\|_{B_{\infty,1}^0} \leq \|\omega(\tau)\|_{B_{\infty,1}^0}. \quad (3.1.5)$$

He then applies (3.1.2) and Theorem 3.1.1 to conclude that

$$\|\nabla v(\tau)\|_{L^\infty} \leq C(1 + \log(\|g(t)\|_{lip}\|g^{-1}(t)\|_{lip}))\|\omega^0\|_{F_{1,q}^2}, \quad (3.1.6)$$

and using this inequality he proves global well-posedness when initial velocity belongs to  $F_{1,q}^3(\mathbb{R}^2)$ .

**Remark 3.1.1.** The space  $F_{1,q}^3(\mathbb{R}^2)$  is equivalent to the space  $W^{3,\tilde{1}}(\mathbb{R}^2)$ , where the  $\tilde{1}$  denotes the Hardy space  $\mathcal{H}^1(\mathbb{R}^2)$  in place of the usual Lebesgue space  $L^1(\mathbb{R}^2)$ . Therefore, Chae's result gives global well-posedness for the Euler equations in this space.

Despite several results in the critical Besov and Triebel-Lizorkin spaces in the plane, many interesting problems remain open for the critical case. In particular, regularity of solutions to the Euler equations with initial velocity in  $W^{s+1,p}(\mathbb{R}^2)$ , for  $sp = 2$ , is not well understood. The main motivation for this chapter is a theorem of Bahouri and Chemin found in [1] which studies this problem. The authors show that a lower bound for the Sobolev exponent of  $\omega(t)$  is determined by the log-Lipschitz norm of  $v(t)$ . They define

$$V(t) = \sup_{|x-y|\leq 1} \frac{|v(t,x) - v(t,y)|}{|x-y|(1 - \log|x-y|)},$$

and they prove the following:

**Theorem 3.1.2.** *Let  $v$  be a solution to (E) such that  $\omega(v^0) = \omega^0 \in L^\infty(\mathbb{R}^2) \cap W^{s,p}(\mathbb{R}^2)$ , for  $sp \leq 2$  and  $s \in (0, 1]$ . Fix  $s' < s$ , and define  $\sigma(s', t) = s' \exp(-\int_0^t V(\tau)d\tau)$ . Then  $\omega(t) \in W^{\sigma(s',t),p}(\mathbb{R}^2)$  for all  $t \in \mathbb{R}$ .*

To prove the theorem, the authors show loss of regularity in the Triebel-Lizorkin space  $F_{p,\infty}^s(\mathbb{R}^2)$ . They use the following definition of  $F_{p,\infty}^s(\mathbb{R}^2)$ :

**Definition 3.1.1.** Let  $s \in (0, 1)$  and  $p \in (1, \infty]$ . The space  $F_{p,\infty}^s(\mathbb{R}^2)$  is the set of all tempered distributions  $u$  in  $L^p(\mathbb{R}^2)$  such that there exists a function

$U$  in  $L^p(\mathbb{R}^2)$  satisfying, for all  $(x, y) \in \mathbb{R}^2 \times \mathbb{R}^2$ ,

$$\frac{|u(x) - u(y)|}{|x - y|^s} \leq U(x) + U(y). \quad (3.1.7)$$

The  $F_{p,\infty}^s$ -norm is given by

$$\|u\|_{F_{p,\infty}^s} = \|u\|_{L^p} + \inf\{\|U\|_{L^p} : U \text{ satisfies (3.1.7)}\}.$$

For a proof that this definition is equivalent to Definition 2.2.4, we refer the reader to [1], Proposition 3.2.

The authors proceed by considering what happens to the Triebel-Lizorkin exponent of  $\omega^0$  under composition with the measure-preserving map  $g(t)$ . Using the fact that  $g(t) - Id$  is Holder continuous with Holder exponent  $\alpha(t) = \exp(-\int_0^t V(\tau)d\tau)$ , they write

$$\begin{aligned} \frac{|\omega^0(g_{t-1}(x)) - \omega^0(g_{t-1}(y))|}{|x - y|^{s\alpha(t)}} &= \frac{|\omega^0(g_{t-1}(x)) - \omega^0(g_{t-1}(y))|}{|g_{t-1}(x) - g_{t-1}(y)|^s} \frac{|g_{t-1}(x) - g_{t-1}(y)|^s}{|x - y|^{s\alpha(t)}} \\ &\leq C(t)^s (U(g_{t-1}(x)) + U(g_{t-1}(y))), \end{aligned} \quad (3.1.8)$$

where  $C(t)$  depends on the Holder norm of  $g(t) - Id$ . Since the flow is measure-preserving, it follows from 3.1.8 and the definition of the  $F_{p,\infty}^s$ -norm that

$$\|\omega^0 \circ g_{t-1}\|_{F_{p,\infty}^{s\alpha(t)}} \leq (C(t)^s + 1) \|\omega^0\|_{F_{p,\infty}^s}.$$

The theorem follows from the embeddings  $W^{s,p}(\mathbb{R}^2) \hookrightarrow F_{p,\infty}^s(\mathbb{R}^2) \hookrightarrow W^{s',p}(\mathbb{R}^2)$  for all  $s' < s$ .

We claim that if we also assume  $\omega^0$  is uniformly continuous, then we can improve the lower bound for loss of regularity to an arbitrarily small amount. We will show that, given  $\epsilon > 0$  arbitrarily small and  $T > 0$  fixed, if  $\omega^0 \in W^{s,p}(\mathbb{R}^2) \cap UC(\mathbb{R}^2)$  with  $sp \leq 2$ ,  $p \in (1, \infty)$ , and  $s \in (0, 2)$ , then  $\omega(t)$  belongs to  $W^{s-\epsilon,p}(\mathbb{R}^2) \cap UC(\mathbb{R}^2)$  for all  $t \in [0, T]$ . As in [1], we study the vorticity equation ( $V$ ) corresponding to the Euler equations.

To prove our result, we show that  $\omega(t)$  belongs to the Besov space  $B_{p,\infty}^{s-\epsilon}$  (see Definition 2.2.3) for all  $t$  in a finite time interval  $[0, T]$ . Our approach is to localize the frequency of the terms of ( $V$ ), which results in a new equation with a commutator term on the right-hand side:

$$\begin{aligned} \partial_t \Delta_q \omega + v \cdot \nabla \Delta_q \omega &= [v \cdot \nabla, \Delta_q] \omega, \\ \Delta_q \omega|_{t=0} &= \Delta_q \omega^0. \end{aligned} \tag{3.1.9}$$

We then prove the necessary estimate on the  $L^p$ -norm of the commutator on the right-hand side of (3.1.9), and apply a Gronwall argument to show that  $\omega(t)$  is in  $B_{p,\infty}^{s-\epsilon}(\mathbb{R}^2)$ .

The main novelty of this proof is that our methods allow us to draw conclusions for  $\omega^0 \in B_{p,\infty}^s(\mathbb{R}^2) \cap UC(\mathbb{R}^2)$  for *all*  $s \in (0, 2)$ . If we restrict our attention to the case  $s \in (0, 1]$ , we can also prove our result by modifying the proof found in [1] (which we have described above) using the following theorem:

**Theorem 3.1.3.** *Let  $v$  be a solution to (E) such that  $\omega(v^0) = \omega^0$  belongs to  $UC(\mathbb{R}^2)$ . Let  $g(t)$  be the measure-preserving homeomorphism in  $\mathbb{R}^2$  satisfying*

$\partial_t g(t, x) = v(t, g(t, x))$ . Given  $\delta > 0$  and  $T > 0$ , it follows that  $\|g(t)^{-1} - Id\|_{C^{1-\delta}}$  belongs to  $L^\infty([0, T])$ .

We devote the next section to a proof of Theorem 3.1.3. We prove the main result in Section 3.4.

### 3.2 A proof of Theorem 3.1.4

An argument using Osgood's Lemma (see, for example, chapter 5 of [6]) implies that the Holder exponent of the flow  $g(t) - Id$  is determined by the log-Lipschitz norm of  $v$ . Precisely,

$$\sup_{|x-y|\leq 1} |g(t)(x) - g(t)(y)| \leq C|x - y|^{\alpha(t)}, \quad (3.2.1)$$

where  $\alpha(t) = \exp(-\int_0^t V(\tau)d\tau)$ , and  $V(t)$  is defined as in section 3.1. One can characterize log-Lipschitz functions using the following inequality (see [1]):

$$C^{-1}\|f\|_{LL} \leq \|S_0 f\|_{L^\infty} + \sup_{q \geq 1} \frac{\|\nabla S_q f\|_{L^\infty}}{q+1} \leq C\|f\|_{LL} \quad (3.2.2)$$

for a constant  $C > 0$ . When computing the Holder exponent of  $g(t) - Id$ , (3.2.1) and (3.2.2) motivate us to study the behavior of the quantity  $\|\nabla S_q v(t)\|_{L^\infty}$ . In this section, we assume  $\omega^0$  belongs to  $UC(\mathbb{R}^2)$ , and we show that, given  $\epsilon > 0$  and  $T > 0$ ,  $\|\nabla S_q v(t)\|_{L^\infty} \leq \epsilon(q+1)$  for sufficiently large  $q$  and for all  $t \in [0, T]$ . We then conclude that  $g(t)$  is locally Holder continuous with Holder exponent arbitrarily close to 1. We begin with the following lemma:

**Lemma 3.2.1.** *Let  $u$  be a uniformly continuous function on  $\mathbb{R}^2$ . Given  $\epsilon > 0$ , there exists an  $N > 0$  such that  $\|\Delta_j u\|_{L^\infty} < \epsilon$  for all  $j > N$ .*

*Proof.* Since  $u$  is bounded and uniformly continuous, it can be approximated uniformly by the sequence  $(S_N u)_{N=1}^\infty$ . Therefore, for  $N$  sufficiently large and for all  $j > N$  we have

$$\begin{aligned} \|\Delta_j u\|_{L^\infty} &\leq \|\Delta_j(S_N u - u)\|_{L^\infty} + \|\Delta_j S_N u\|_{L^\infty} \\ &\leq C\|S_N u - u\|_{L^\infty} < \epsilon. \end{aligned}$$

This completes the proof. □

Observe that by Remark 2.3.1 in Section 2.3.2, the solution to (V) satisfies  $\omega(t, x) = \omega^0(g(t)^{-1}(x))$ . Therefore, if we assume that  $\omega^0$  is uniformly continuous on  $\mathbb{R}^2$ , then  $\omega(t)$  is uniformly continuous on  $\mathbb{R}^2$  for all  $t$  in  $\mathbb{R}$ . We now apply Lemma 3.2.1 to  $\omega(t)$  and conclude that, for fixed  $t$ , given  $\epsilon > 0$ , there exists  $N_t$  such that  $\sup_{j > N_t} \|\Delta_j \omega(t)\|_{L^\infty} < \epsilon$ . In what follows, we need  $N_t$  to be time independent. We therefore prove the following lemma:

**Lemma 3.2.2.** *Let  $v$  be a solution to (E) such that  $\omega(v^0) = \omega^0$  belongs to  $UC(\mathbb{R}^2)$ . Given  $\epsilon > 0$  and  $T > 0$ , there exists an  $N = N(T, \epsilon)$  such that  $\sup_{j > N} \|\Delta_j \omega(t)\|_{L^\infty} < \epsilon$  for all  $t \in [0, T]$ .*

*Proof.* Observe that, for  $x_1$  and  $x_2$  in  $\mathbb{R}^2$  satisfying  $g(t_1, x_1) = g(t_2, x_2) = x$ ,

we have,

$$\begin{aligned}
|x_1 - x_2| &\leq |t_1 - t_2| \sup_{\tau \in [t_1, t_2]} \|v(\tau)\|_{L^\infty} \\
&\leq C|t_1 - t_2| \|\omega^0\|_{L^p} e^{C_1 t_2 \|\omega^0\|_{L^p}} \\
&\leq C|t_1 - t_2| \|\omega^0\|_{L^p} e^{C_1 T \|\omega^0\|_{L^p}},
\end{aligned}$$

where we used Lemma 2.3.11. Therefore, by uniform continuity of  $\omega^0$ , given  $\epsilon > 0$  there exists  $\delta = \delta(T) > 0$  such that for  $|t_1 - t_2| < \delta$  and for all  $x \in \mathbb{R}^2$ ,

$$|\omega(t_1, x) - \omega(t_2, x)| = |\omega^0(x_1) - \omega^0(x_2)| < \frac{\epsilon}{2}.$$

We now break  $[0, T]$  into intervals of length less than or equal to  $\delta$ :  $[0, t_1], [t_1, t_2], \dots, [t_{M-1}, T]$ . We observe that, for each  $t \in [0, T]$ , there exists an  $i$ ,  $1 \leq i \leq M$ , such that  $|t - t_i| < \delta$ , and therefore  $\|\Delta_j \omega(t)\|_{L^\infty} \leq \frac{\epsilon}{2} + \|\Delta_j \omega(t_i)\|_{L^\infty}$ . Furthermore, given  $\omega(t_i)$ , there exists an  $N_{t_i}$  such that  $\sup_{j > N_{t_i}} \|\Delta_j \omega(t_i)\|_{L^\infty} < \frac{\epsilon}{2}$ . Let  $N = \max\{N_{t_1}, \dots, N_{t_M}\}$ . Then, for all  $t \in [0, T]$ ,

$$\sup_{j > N} \|\Delta_j \omega(t)\|_{L^\infty} \leq \frac{\epsilon}{2} + \sup_{j > N} \|\Delta_j \omega(t_i)\|_{L^\infty} \leq \epsilon.$$

This completes the proof.  $\square$

We use Lemma 2.3.8 to bound  $\|\Delta_j \nabla v\|_{L^\infty}$  by  $C\|\Delta_j \omega\|_{L^\infty}$  if  $j \geq 0$ , and we bound  $\|\Delta_{-1} \nabla v(t)\|_{L^\infty}$  with  $C\|\omega^0\|_{L^p}$  for  $p \in (1, \infty)$  using Bernstein's inequality. From Lemma 3.2.2, we conclude that, for  $N$  sufficiently large, and for all  $t \in [0, T]$ ,

$$\|S_N \nabla v(t)\|_{L^\infty} \leq \epsilon(N + 1). \quad (3.2.3)$$

**Remark 3.2.1.** One can also prove (3.2.3) using Calderon-Zygmund theory. As in Lemma 3.2.1, since  $\omega$  is bounded and uniformly continuous, it can be approximated uniformly by the sequence of functions  $(S_N\omega)_{N=1}^\infty$ . Because  $\omega$  is  $L^p$  integrable for some  $p \in (1, \infty)$ , the functions in this sequence vanish at infinity; therefore,  $\omega$  vanishes at infinity. By the theory of Calderon-Zygmund operators, since  $\omega$  is continuous, in  $L^p(\mathbb{R}^2)$  for some  $p \in (1, \infty)$ , and vanishes at infinity,  $\nabla v$  belongs to the space  $vmo(\mathbb{R}^2)$  (see, for example, [16]). This implies by Definition 2.2.8 that there exists a sequence of compactly supported  $C^\infty$  functions  $(u_N)_{N=1}^\infty$  converging to  $\nabla v$  in  $bmo(\mathbb{R}^2)$ . But  $bmo(\mathbb{R}^2)$  imbeds into the Zygmund space  $C_*^0(\mathbb{R}^2)$  by Proposition 2.2.1; thus the sequence converges to  $\nabla v$  in  $C_*^0(\mathbb{R}^2)$ , and we can write

$$\|\Delta_j \nabla v\|_{L^\infty} \leq \|\Delta_j(\nabla v - u_n)\|_{L^\infty} + \|\Delta_j u_n\|_{L^\infty}. \quad (3.2.4)$$

We choose  $n$  so that the first term on the right hand side of (3.2.4) is small for all  $j$ . Given this  $n$ , we choose  $j$  large enough to make the second term on the right hand side of (3.2.4) as small as we would like. This gives Lemma 3.2.1, and, following the argument above, implies (3.2.3).

We now use (3.2.3) to compute the Holder exponent for the flow corresponding to the velocity of a fluid with initial vorticity in  $UC(\mathbb{R}^2)$ , which will complete the proof of Theorem 3.1.3. We show that, given  $\epsilon > 0$  and  $T > 0$ ,  $g(t)^{-1} - Id$  belongs to  $C^{\sigma(t)}(\mathbb{R}^2)$  for all  $t \in [0, T]$ , where  $\sigma(t) = e^{-Ct\epsilon}$ , and  $C$  is an absolute constant.

Fix  $\epsilon > 0$ . Write  $v = v_{1,N} + v_{2,N}$ , where  $v_{1,N} = S_{N-1}v$ , and  $v_{2,N} = (Id - S_{N-1})v$ . By (3.2.3), it follows that  $|v_{1,N}(t, x) - v_{1,N}(t, y)| \leq \|\nabla S_{N-1}v(t)\|_{L^\infty} |x - y| \leq CN\epsilon |x - y|$  for large enough  $N$ . Similarly, for large  $N$  we can conclude that  $|v_{2,N}(t, x) - v_{2,N}(t, y)| \leq C \sum_{j=N-1}^{\infty} 2^{-j} \|\Delta_j \nabla v(t)\|_{L^\infty} \leq C2^{-N}\epsilon$ . Letting  $N = -\log_2 |x - y|$ , we have that for  $|x - y|$  sufficiently small,

$$\begin{aligned} |v(t, x) - v(t, y)| &\leq |(v_{1,N}(t, x) + v_{2,N}(t, x)) - (v_{1,N}(t, y) + v_{2,N}(t, y))| \\ &\leq C(\epsilon(-\log_2 |x - y|)|x - y| + \epsilon|x - y|) \\ &\leq C\epsilon(1 - \log_2 |x - y|)|x - y|. \end{aligned} \tag{3.2.5}$$

We now use (3.2.5) and Osgood's Lemma (see Lemma 2.3.4) to compute properties of the flow. We write

$$|v(t, g(t, x)) - v(t, g(t, y))| \leq C\epsilon(1 - \log_2 |g(t, x) - g(t, y)|)|g(t, x) - g(t, y)|$$

whenever  $|g(t, x) - g(t, y)| < \delta$ . From (2.3.6), we see that  $|g(t, x) - g(t, y)|$  is bounded above by

$$|x - y| + \int_0^t C\epsilon(1 - \log_2 |g(\tau, x) - g(\tau, y)|)|g(\tau, x) - g(\tau, y)| d\tau.$$

By Osgood's Lemma, we conclude that

$$-\log(1 - \log |g(t, x) - g(t, y)|) + \log(1 - \log |x - y|) \leq Ct\epsilon.$$

Taking the exponential twice, we get

$$\frac{|g(t, x) - g(t, y)|}{|x - y|^{e^{-Ct\epsilon}}} \leq e^{1 - e^{-Ct\epsilon}} \leq e$$

whenever  $|x - y| < \tilde{\delta}$ , where  $\tilde{\delta} = e^{1 - e^{-Ct\epsilon}}$ . This gives

$$\frac{|(g(t, x) - x) - (g(t, y) - y)|}{|x - y|^{e^{-Ct\epsilon}}} \leq e + 1.$$

In the case  $|x - y| \geq \tilde{\delta}$ , we have

$$\frac{|(g(t, x) - x) - (g(t, y) - y)|}{|x - y|^{e^{-Ct\epsilon}}} \leq 2\tilde{\delta}^{-e^{-Ct\epsilon}} \|g(t) - Id\|_{L^\infty}.$$

To see that  $\|g(t)^{-1} - Id\|_{C^{e^{-Ct\epsilon}}} \in L_{loc}^\infty(\mathbb{R}^+)$ , we observe that

$$\begin{aligned} \sup_{x, y \in \mathbb{R}^2} \frac{|(g(t, x) - x) - (g(t, y) - y)|}{|x - y|^{e^{-Ct\epsilon}}} &\leq (e + 1) + 2\tilde{\delta}^{-e^{-Ct\epsilon}} \left\| \int_0^t v(\tau, g(\tau, \cdot)) d\tau \right\|_{L^\infty} \\ &\leq (e + 1) + 2\tilde{\delta}^{-1} T (\|\omega^0\|_{L^p} + \|\omega^0\|_{L^\infty}) \end{aligned}$$

for all  $t \in [0, T]$ . This completes the proof of Theorem 3.1.3.

### 3.3 Paradifferential estimates for the vorticity equation

In this section, we consider an initial value problem for the vorticity equation corresponding to the two-dimensional Euler equations. Recall that in dimension two, the vorticity equation is given by

$$(V) \quad \begin{cases} \partial_t \omega + u \cdot \nabla \omega = 0, \\ \omega|_{t=0} = \omega^0. \end{cases}$$

Note that, if  $\omega$  satisfies (V), then  $\Delta_q \omega$  satisfies the following equation:

$$(V^*) \quad \begin{cases} \partial_t \Delta_q \omega + v \cdot \nabla \Delta_q \omega = [v \cdot \nabla, \Delta_q] \omega, \\ \Delta_q \omega|_{t=0} = \Delta_q \omega^0. \end{cases}$$

We want to prove the following estimate:

**Proposition 3.3.1.** *Let  $p \in (1, \infty)$  and  $\sigma > 0$  be fixed. Then there exist two positive constants  $C_1(\sigma)$  and  $C_2$  such that*

$$\|[v \cdot \nabla, \Delta_q] \omega\|_{L^p} \leq C_1(\sigma) (C_2 + \|S_{q-1} \nabla v\|_{L^\infty}) 2^{-q\sigma} \|\omega\|_{B_{p, \infty}^\sigma}.$$

**Remark 3.3.1.** To simplify notation, we suppress the time variable in the statement and proof of Proposition 3.3.1 (time is fixed throughout the proof). However, we will reintroduce the time variable when we use Proposition 3.3.1 to prove the main theorem in Section 3.4. Therefore, it is important to emphasize that the constants  $C_1$  and  $C_2$  will depend on time in Section 3.4. In the proof of Proposition 3.3.1, we find that  $C_1$  depends on  $\sigma$ , and in Section 3.4 we let  $\sigma$  vary with time. We also find in the proof of Proposition 3.3.1 that  $C_2 = C\|\omega^0\|_{L^p \cap L^\infty} e^{Ct\|\omega^0\|_{L^p}}$ , where  $C$  is an absolute constant, and  $p \in (2, \infty)$ . This dependence arises from bounding  $\|v(t)\|_{L^\infty}$  as in Lemma 2.3.11. (If in fact  $\omega^0$  belongs to  $L^p(\mathbb{R}^2)$  for some  $p < 2$ , then it is possible to bound  $\|v(t)\|_{L^\infty}$  by a constant which depends only on initial vorticity and does not depend on time.)

*Proof.* We consider the cases  $q \geq 4$  and  $q < 4$  separately. We first assume  $q \geq 4$  and use Bony's decomposition to write

$$[v \cdot \nabla, \Delta_q] \omega = \sum_{j=1}^2 [T_{v^j} \partial_j, \Delta_q] \omega + [T_{\partial_j} v^j, \Delta_q] \omega + [\partial_j R(v^j, \cdot), \Delta_q] \omega.$$

We address each piece of the sum separately. We start with  $[T_{v^j} \partial_j, \Delta_q] \omega$ . Write

$$[T_{v^j} \partial_j, \Delta_q] \omega = \sum_{q'=q-4}^{q+4} [S_{q'-1}(v^j), \Delta_q] \Delta_{q'} \partial_j \omega.$$

Letting  $u = \Delta_{q'} \partial_j \omega$ , letting  $h = \check{\phi}$  (recall we are assuming  $q \geq 4$  here), and

keeping in mind that  $|q' - q| \leq 4$ , we have

$$\begin{aligned}
& \| [S_{q'-1}(v^j), \Delta_q] u \|_{L^p} \\
&= \left\| \int_{\mathbb{R}^2} h(y) (S_{q'-1}(v^j)(x - 2^{-q}y) - S_{q'-1}(v^j)(x)) u(x - 2^{-q}y) dy \right\|_{L^p} \\
&\leq C \| S_{q'-1} \nabla v \|_{L^\infty} 2^{-q} \| u \|_{L^p} \\
&\leq C 2^{4\sigma} \| S_{q'-1} \nabla v \|_{L^\infty} 2^{-q\sigma} \| \omega \|_{B_{p,\infty}^\sigma},
\end{aligned}$$

where we used the fact that  $h \in S$  and therefore  $zh(z)$  is integrable, as well as Bernstein's inequality. We now sum over  $q'$  to get

$$\begin{aligned}
\| [T_{v^j} \partial_j, \Delta_q] \omega \|_{L^p} &\leq C 2^{4\sigma} 2^{-q\sigma} \| \omega \|_{B_{p,\infty}^\sigma} \sum_{q'=q-4}^{q+4} \| S_{q'-1} \nabla v \|_{L^\infty} \\
&\leq C 2^{4\sigma} 2^{-q\sigma} \| \omega \|_{B_{p,\infty}^\sigma} (\| S_{q-1} \nabla v \|_{L^\infty} + \| \nabla v \|_{C_*^0}).
\end{aligned} \tag{3.3.1}$$

We now consider  $[T_{\partial_j \cdot v^j}, \Delta_q] \omega$ . To bound  $\| T_{\partial_j \Delta_q \omega} v^j \|_{L^p}$ , we use Bernstein's inequality and our assumption that  $q \geq 4$ , as well as properties of our partition of unity, to write

$$\begin{aligned}
\| T_{\partial_j \Delta_q \omega} v^j \|_{L^p} &\leq \sum_{q'=q}^{\infty} C 2^q 2^{-q'} \| S_{q'-1} \Delta_q \omega \|_{L^p} \| \Delta_{q'} \nabla v \|_{L^\infty} \\
&\leq C \| \Delta_q \omega \|_{L^p} \sup_{q' \geq q} \| \Delta_{q'} \nabla v \|_{L^\infty} \\
&\leq C \| \nabla v \|_{C_*^0} 2^{-q\sigma} \| \omega \|_{B_{p,\infty}^\sigma}.
\end{aligned} \tag{3.3.2}$$

Furthermore, since the Fourier support of  $S_{q'-1} \partial_j \omega \Delta_{q'} v^j$  is contained in an

annulus with inner and outer radius  $C2^{q'}$  and  $\tilde{C}2^{q'}$  respectively, we can write

$$\begin{aligned}
\|\Delta_q(T_{\partial_j\omega}v^j)\|_{L^p} &\leq \sum_{q'=q-4}^{q+4} \|S_{q'-1}\partial_j\omega\|_{L^\infty} \|\Delta_{q'}v\|_{L^p} \\
&\leq \sum_{q'=q-4}^{q+4} C2^{q'}2^{-q'} \|S_{q'-1}\omega\|_{L^\infty} \|\Delta_{q'}\nabla v\|_{L^p} \\
&\leq \sum_{q'=q-4}^{q+4} C \|S_{q'-1}\nabla v\|_{L^\infty} \|\Delta_{q'}\omega\|_{L^p} \\
&\leq C2^{4\sigma} (\|S_{q-1}\nabla v\|_{L^\infty} + \|\nabla v\|_{C_*^0}) 2^{-q\sigma} \|\omega\|_{B_{p,\infty}^\sigma}.
\end{aligned} \tag{3.3.3}$$

Once again, we used Bernstein's inequality in the second inequality. It is in this inequality that our assumption that  $q \geq 4$  is necessary. For the third inequality, we used Lemma 2.3.8. Combining (3.3.2) and (3.3.3), we see that

$$\|[T_{\partial_j}v^j, \Delta_q]\omega\|_{L^p} \leq C2^{4\sigma} (\|S_{q-1}\nabla v^j\|_{L^\infty} + \|\nabla v\|_{C_*^0}) 2^{-q\sigma} \|\omega\|_{B_{p,\infty}^\sigma} \tag{3.3.4}$$

for  $q \geq 4$ .

We now study the remainder term,  $[\partial_j R(v^j, \cdot), \Delta_q]\omega$ . We need the following lemma:

**Lemma 3.3.2.** *If  $s + \sigma > 0$ , then  $\|R(a, b)\|_{B_{p,\infty}^{s+\sigma}} \leq C(s, \sigma) \|a\|_{B_{\infty,\infty}^s} \|b\|_{B_{p,\infty}^\sigma}$ , where  $C(s, \sigma) = C2^{M\sigma} (\frac{1}{1-(\frac{1}{2})^{s+\sigma}} + 2^{N(s+\sigma)} + \dots + 2^{1(s+\sigma)})$  for fixed positive integers  $M$  and  $N$ .*

*Proof.* For a proof of this lemma, see [22]. □

To handle the remainder term, we will consider low and high frequencies of  $v^j$  separately. We begin with  $[\partial_j R((Id - \Delta_{-1})v^j, \cdot), \Delta_q]\omega$ . Using Lemma

3.3.2 with  $s = 1$ , Bernstein's inequality, and the fact that the Fourier transform of  $(Id - \Delta_{-1})\nabla v$  vanishes in a neighborhood of the origin, we write

$$\begin{aligned}
\|\Delta_q \partial_j R((Id - \Delta_{-1})v^j, \omega)\|_{L^p} &\leq 2^{-q\sigma} \|R((Id - \Delta_{-1})v^j, \omega)\|_{B_{p,\infty}^{\sigma+1}} \\
&\leq C(\sigma) 2^{-q\sigma} \|(Id - \Delta_{-1})v\|_{B_{\infty,\infty}^1} \|\omega\|_{B_{p,\infty}^\sigma} \\
&\leq C(\sigma) 2^{-q\sigma} \|\nabla v\|_{C_*^0} \|\omega\|_{B_{p,\infty}^\sigma}.
\end{aligned} \tag{3.3.5}$$

Here  $C(\sigma) = C(1, \sigma)$  from Lemma 3.3.2. To bound  $\|\partial_j R((Id - \Delta_{-1})v^j, \Delta_q \omega)\|_{L^p}$ , note that

$$\begin{aligned}
\|\partial_j R((Id - \Delta_{-1})v^j, \Delta_q \omega)\|_{L^p} &\leq \sum_{\substack{q', q'' \\ |q' - q''| \leq 1 \\ |q' - q| \leq 1}} 2^q \|\Delta_{q''}(Id - \Delta_{-1})v\|_{L^\infty} \|\Delta_{q'} \Delta_q \omega\|_{L^p} \\
&\leq C \|\nabla v\|_{C_*^0} 2^{-q\sigma} \|\omega\|_{B_{p,\infty}^\sigma}.
\end{aligned} \tag{3.3.6}$$

In the first inequality above, we used the fact that the support of the Fourier transform of  $\Delta_{q''}(Id - \Delta_{-1})v \Delta_{q'} \Delta_q \omega$  is contained in a ball with radius  $C2^q$ , along with Bernstein's inequality, to get the factor  $2^q$ . In the second inequality, we used the inequality  $\|(Id - \Delta_{-1})v\|_{B_{\infty,\infty}^1} \leq C \|\nabla v\|_{C_*^0}$ . We now combine (3.3.6) with (3.3.5) to conclude that

$$\|\partial_j R((Id - \Delta_{-1})v^j, \cdot), \Delta_q \omega\|_{L^p} \leq C(\sigma) \|\nabla v\|_{C_*^0} 2^{-q\sigma} \|\omega\|_{B_{p,\infty}^\sigma}. \tag{3.3.7}$$

We now estimate  $\|[\partial_j R(\Delta_{-1}v^j, \cdot), \Delta_q \omega]\|_{L^p}$ . Using the definition of the remainder operator, as well as the properties of our partition of unity, we write

$$\begin{aligned}
[\partial_j R(\Delta_{-1}v^j, \cdot), \Delta_q \omega] &= \partial_j R(\Delta_{-1}v^j, \Delta_q \omega) - \Delta_q(\partial_j R(\Delta_{-1}v^j, \omega)) \\
&= \partial_j \left( \sum_{\substack{i,k \\ |i-k| \leq 1}} \Delta_k \Delta_{-1} v^j \Delta_i \Delta_q \omega \right) - \Delta_q \partial_j \left( \sum_{\substack{i,k \\ |i-k| \leq 1}} \Delta_k \Delta_{-1} v^j \Delta_i \omega \right).
\end{aligned}$$

We begin by estimating  $\Delta_q \partial_j (\sum_{\substack{i,k \\ |i-k| \leq 1}} \Delta_k \Delta_{-1} v^j \Delta_i \omega)$ . We first reintroduce the sum over  $j$ , allowing us to use the fact that  $\operatorname{div} v = 0$  to move  $\partial_j$  inside the parentheses and differentiate  $\omega$ . This, along with properties of our partition of unity, gives

$$\begin{aligned} & \sum_{j=1}^2 \|\Delta_q \partial_j (\sum_{\substack{i,k \\ |i-k| \leq 1}} \Delta_k \Delta_{-1} v^j \Delta_i \omega)\|_{L^p} \\ & \leq C \sum_{j=1}^2 \sum_{l=-1}^1 \sum_{k=-1}^0 \|\Delta_q (\Delta_k \Delta_{-1} v^j \Delta_{k-l} \partial_j \omega)\|_{L^p}. \end{aligned} \quad (3.3.8)$$

Note that in the second line of (3.3.8), the Fourier support of  $\Delta_k \Delta_{-1} v^j \Delta_{k-l} \partial_j \omega$  is contained in a ball with radius  $C2^k$ . Therefore, the sum in the second line is 0 if  $q \geq k + M$ , for a constant  $M$ . Furthermore,  $k \leq 0$ . Therefore, we are only considering  $q \leq M$ . We then write

$$\begin{aligned} & \sum_{j=1}^2 \sum_{l=-1}^1 \sum_{k=-1}^0 \|\Delta_q (\Delta_k \Delta_{-1} v^j \Delta_{k-l} \partial_j \omega)\|_{L^p} \\ & \leq C 2^{M\sigma} \|v\|_{L^\infty} 2^{-q\sigma} \|\omega\|_{B_{p,\infty}^\sigma}. \end{aligned} \quad (3.3.9)$$

To bound  $\|\partial_j (\sum_{\substack{i,k \\ |i-k| \leq 1}} \Delta_k \Delta_{-1} v^j \Delta_i \Delta_q \omega)\|_{L^p}$ , we use the fact that the Fourier transform of  $\Delta_{-1} v^j$  is supported in the neighborhood of the origin, and we recognize that  $\operatorname{div} v = 0$  allows us to move  $\partial_j$  inside the parentheses. Therefore,

$$\begin{aligned} & \sum_{j=1}^2 \|\partial_j (\sum_{\substack{i,k \\ |i-k| \leq 1}} \Delta_k \Delta_{-1} v^j \Delta_i \Delta_q \omega)\|_{L^p} \\ & \leq C \sum_{j=1}^2 \sum_{l=-1}^1 \sum_{k=-1}^0 \|\Delta_k \Delta_{-1} v\|_{L^\infty} 2^{k-l} \|\Delta_{k-l} \Delta_q \omega\|_{L^p} \\ & \leq C \|v\|_{L^\infty} 2^{-q\sigma} \|\omega\|_{B_{p,\infty}^\sigma}, \end{aligned} \quad (3.3.10)$$

where we used Bernstein's inequality and Holder's inequality to get the first inequality in (3.3.10). We now combine (3.3.8) through (3.3.10) to conclude that

$$\|[\partial_j R(\Delta_{-1} v^j, \cdot), \Delta_q] \omega\|_{L^p} \leq C 2^{M\sigma} \|v\|_{L^\infty} 2^{-q\sigma} \|\omega\|_{B_{p,\infty}^\sigma}. \quad (3.3.11)$$

Combining (3.3.1), (3.3.4), (3.3.7), and (3.3.11), we conclude that for  $q \geq 4$ ,

$$\begin{aligned} \|[v \cdot \nabla, \Delta_q] \omega\|_{L^p} &\leq C(2^{M\sigma} + C(\sigma)) (\|S_{q-1} \nabla v^j\|_{L^\infty} \\ &\quad + \|\nabla v\|_{C_*^0} + \|v\|_{L^\infty}) 2^{-q\sigma} \|\omega\|_{B_{p,\infty}^\sigma}. \end{aligned}$$

To complete the proof for the case  $q \geq 4$ , we use Lemma 2.3.11 to bound  $\|\nabla v\|_{C_*^0} + \|v\|_{L^\infty}$  by  $C_2$ , where  $C_2$  is defined as in Remark 3.3.1. This completes the proof for the case  $q \geq 4$ .

For the case  $q < 4$ , write:

$$[v \cdot \nabla, \Delta_q] \omega = v \cdot \nabla \Delta_q \omega - \Delta_q (v \cdot \nabla \omega). \quad (3.3.12)$$

Keeping in mind that  $q \leq 3$ , it is easy to see that

$$\|v \cdot \nabla \Delta_q \omega\|_{L^p} \leq C_2 2^{-q\sigma} \|\omega\|_{B_{p,\infty}^\sigma}, \quad (3.3.13)$$

where again we used Lemma 2.3.11 to bound  $\|v\|_{L^\infty}$ . We now write the second term of (3.3.12) as

$$\Delta_q (v \cdot \nabla \omega) = \sum_{j=1}^2 \Delta_q (T_{v^j} \partial_j \omega + T_{\partial_j \omega} v^j + R(v^j, \partial_j \omega)). \quad (3.3.14)$$

We successfully bounded the  $L^p$ -norm of the remainder term of (3.3.14) in (3.3.5), (3.3.8), and (3.3.9) of the proof for the  $q \geq 4$  case (note that (3.3.5),

(3.3.8), and (3.3.9) hold for all  $q$ ). Therefore, we are only concerned with  $\sum_{j=1}^2 \Delta_q(T_{v^j} \partial_j \omega + T_{\partial_j \omega} v^j)$ . Using the fact that  $S_{q'-1} v^j \Delta_{q'} \partial_j \omega$  has Fourier support in an annulus with inner radius  $C2^{q'}$  and outer radius  $\tilde{C}2^{q'}$ , and, once again, keeping in mind that  $q \leq 3$ , we have

$$\begin{aligned} \|\Delta_q(T_{v^j} \partial_j \omega)\|_{L^p} &\leq \sum_{q'=q-4}^{q+4} C \|S_{q'-1} v\|_{L^\infty} 2^{q'} \|\Delta_{q'} \omega\|_{L^p} \\ &\leq C_2 2^{-q\sigma} \|\omega\|_{B_{p,\infty}^\sigma}. \end{aligned} \quad (3.3.15)$$

Furthermore, since  $q \leq 3$ , we write

$$\begin{aligned} \|\Delta_q(T_{\partial_j \omega} v^j)\|_{L^p} &\leq \sum_{q'=q-4}^{q+4} \|S_{q'-1} \partial_j \omega \Delta_{q'} v^j\|_{L^p} \\ &\leq C \sum_{q'=q-4}^{q+4} \sum_{k=-1}^{q'-2} 2^{-k\sigma} \|\omega\|_{B_{p,\infty}^\sigma} \|v\|_{L^\infty} \\ &\leq C_2 2^{3\sigma} 2^{-q\sigma} \|\omega\|_{B_{p,\infty}^\sigma}. \end{aligned}$$

This completes the proof of the case  $q < 4$ , and therefore completes the proof of the estimate for all  $q$ . We conclude that, for all  $q$ ,

$$\|[v \cdot \nabla, \Delta_q] \omega\|_{L^p} \leq C_1(\sigma) (\|S_{q-1} \nabla v^j\|_{L^\infty} + C_2) 2^{-q\sigma} \|\omega\|_{B_{p,\infty}^\sigma}.$$

□

If we define  $h(t, x) = g(t)^{-1}(x) - x$ , then  $h$  satisfies the following:

$$\begin{aligned} \partial_t h + v \cdot \nabla h + v &= 0, \\ h|_{t=0} &= 0. \end{aligned} \quad (3.3.16)$$

Since  $h$  satisfies (3.3.16), it also satisfies

$$\begin{aligned} \partial_t \Delta_q h + v \cdot \nabla \Delta_q h &= -\Delta_q v + [v \cdot \nabla, \Delta_q] h, \\ \Delta_q h|_{t=0} &= 0. \end{aligned} \quad (3.3.17)$$

This motivates us to prove a similar commutator estimate with  $h$  in place of  $\omega$ . We prove that the following estimate holds:

**Proposition 3.3.3.** *Let  $p \in (1, \infty)$ ,  $\delta > 0$ , and  $\sigma > 0$  be fixed. There exist two positive constants  $C_1(\sigma)$  and  $C_2$  such that*

$$\begin{aligned} \|[v \cdot \nabla, \Delta_q]h\|_{L^p} &\leq C_1(\sigma)(C_2 + \|S_{q-1}\nabla v\|_{L^\infty})2^{-q\sigma}\|h\|_{B_{p,\infty}^\sigma} \\ &\quad + \sum_{q'=q-4}^{q+4} C(\delta)2^{q'\delta}\|\Delta_{q'}v\|_{L^p}\|h\|_{C^{1-\delta}}. \end{aligned}$$

*Proof.* The proof of Proposition 3.3.3 is identical to the proof of Proposition 3.3.1 with  $h$  in place of  $\omega$  for every term except  $\Delta_q(T_{\partial_j h}v^j)$ . Therefore, we restrict our attention to this term. This portion of the proof will result in the second piece on the right-hand side in Proposition 3.3.3.

Note that, in the proof of Proposition 3.3.1, we use the assumption that  $q \geq 4$  only when bounding  $\|\Delta_q(T_{\partial_j \omega}v^j)\|_{L^p}$ . For all other terms,  $q \geq 0$  suffices. This observation, combined with the fact that we will only need to assume  $q \geq 0$  to bound  $\|\Delta_q(T_{\partial_j h}v^j)\|_{L^p}$ , leads us to consider the cases  $q = -1$  and  $q \geq 0$  separately.

We first assume  $q \geq 0$ , and we write

$$\Delta_q(T_{\partial_j h}v^j) = \Delta_q\left(\sum_{q'=1}^{\infty} S_{q'-1}\partial_j h \Delta_{q'}v^j\right). \quad (3.3.18)$$

Using the fact that  $S_{q'-1}\partial_j h \Delta_{q'}v^j$  has Fourier support in an annulus with inner radius  $C2^{q'}$  and outer radius  $\tilde{C}2^{q'}$ , we apply Bernstein's inequality and Holder's

inequality to get

$$\begin{aligned} \|\Delta_q(T_{\partial_j h} v^j)\|_{L^p} &\leq \sum_{q'=q-4}^{q+4} \sum_{k=-1}^{q'-2} 2^{k\delta} 2^{k(1-\delta)} \|\Delta_k h\|_{L^\infty} \|\Delta_{q'} v\|_{L^p} \\ &\leq \sum_{q'=q-4}^{q+4} C(\delta) 2^{q'\delta} \|h\|_{C^{1-\delta}} \|\Delta_{q'} v\|_{L^p}. \end{aligned} \quad (3.3.19)$$

We now consider the case  $q = -1$ . As in the proof of Proposition 3.3.1 when assuming  $q < 4$ , we begin by writing:

$$[v \cdot \nabla, \Delta_q]h = v \cdot \nabla \Delta_q h - \Delta_q(v \cdot \nabla h). \quad (3.3.20)$$

For the first term of (3.3.20), we use the fact that  $q = -1$  to get

$$\|v \cdot \nabla \Delta_q h\|_{L^p} \leq C(\|\omega^0\|_{L_0^p} + \|\omega^0\|_{L^\infty}) 2^{-q\sigma} \|h\|_{B_{p,\infty}^\sigma}, \quad (3.3.21)$$

for  $p \in (1, 2)$ . The second term of (3.3.20) can be written as

$$\Delta_q(v \cdot \nabla h) = \sum_{j=1}^2 \Delta_q(T_{v^j} \partial_j h + T_{\partial_j h} v^j + R(v^j, \partial_j h)). \quad (3.3.22)$$

The proof of the bound on the  $L^p$ -norm of the remainder term in (3.3.22) is identical to the proofs of (3.3.5), (3.3.8), and (3.3.9), (which hold for all  $q$ ), with  $h$  in place of  $\omega$ . Furthermore, we handled the  $L^p$ -norm of  $\Delta_q(T_{\partial_j h} v^j)$  (see (3.3.19), which also holds for all  $q$ ). Therefore, in (3.3.22), we are only concerned with  $\Delta_q(T_{v^j} \partial_j h)$ . For this term, we refer the reader to the proof for  $\omega$  with  $q < 4$ , given in (3.3.15). This completes the proof of the case  $q = -1$ , and therefore completes the proof of the estimate for all  $q$ .  $\square$

### 3.4 Statement and proof of the main results

In this section we study an initial value problem for ideal incompressible fluids with uniformly continuous vorticity in a Besov space.

#### 3.4.1 Besov regularity of the vorticity

We prove the following theorem:

**Theorem 3.4.1.** *Let  $v^0 \in B_{p,\infty}^{s+1}(\mathbb{R}^2)$ ,  $\operatorname{div} v^0 = 0$ , and let  $\omega(v^0) = \omega^0 \in UC(\mathbb{R}^2)$ , where  $sp \leq 2$ ,  $s \in (0, 2)$ , and  $p \in (1, \infty)$ . Let  $\epsilon > 0$ . There exists a unique solution to (E) such that  $\|v(t)\|_{B_{p,\infty}^{s+1-\epsilon}}$  belongs to  $L_{loc}^\infty(\mathbb{R}^+)$ .*

**Remark 3.4.1.** Uniqueness in Theorem 3.4.1 follows from [23].

*Proof.* Our approach is as follows: we fix  $\epsilon' > 0$  and  $T > 0$ , and we define  $\sigma(t) = s \exp\{-\frac{C}{s}\epsilon't\}$ , where  $C$  is an absolute constant. We then show that  $\|v(t)\|_{B_{p,\infty}^{\sigma(t)+1}} \in L^\infty([0, T])$ . Letting  $\epsilon = s - s \exp\{-\frac{C}{s}\epsilon'T\}$ , we make  $\epsilon$  as small as we would like by our choice of  $\epsilon'$ .

We first prove the theorem on a sufficiently small time interval  $[t_0, t]$ . We then use a bootstrapping argument to show that the theorem holds on any finite time interval  $[0, T]$ . From  $(V^*)$  and Proposition 3.3.1, it follows that

$$\begin{aligned} \|\Delta_q \omega(t)\|_{L^p} &\leq \|\Delta_q \omega(t_0)\|_{L^p} \\ &+ \int_{t_0}^t C_1(\sigma(\tau))(C e^{C\tau} + \|S_{q-1} \nabla v(\tau)\|_{L^\infty}) 2^{-q\sigma(\tau)} \|\omega(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} d\tau. \end{aligned}$$

We used Remark 3.3.1 to replace  $C_2(t)$  with  $C e^{Ct}$  in the above inequality, where  $C$  depends only on the initial vorticity. Moreover, we see from the proof

of Proposition 3.3.1 that  $C_1(\sigma(t))$  can be bounded by an absolute constant for all  $\sigma(t) \in (0, 2)$ . Therefore, for the remainder of the proof, we drop the dependence of  $C_1$  on  $\sigma(t)$ . We multiply both sides of the equation by  $2^{q\sigma(t)}$  and take the supremum over  $q$  to get

$$\begin{aligned} \|\omega(t)\|_{B_{p,\infty}^{\sigma(t)}} &\leq \|\omega(t_0)\|_{B_{p,\infty}^{\sigma(t_0)}} \\ &+ \sup_q \left\{ \int_{t_0}^t C_1(e^{C\tau} + \|S_{q-1}\nabla v(\tau)\|_{L^\infty}) 2^{q\sigma(t)-q\sigma(\tau)} \|\omega(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} d\tau \right\}. \end{aligned}$$

We now show that the supremum over  $q$  on the right-hand side is finite. We claim that the loss of regularity in the Besov exponent, resulting in the term  $2^{q(\sigma(t)-\sigma(\tau))}$ , is enough to combat the growth of  $\|S_{q-1}\nabla v(\tau)\|_{L^\infty}$ .

When taking the supremum over  $q$  of the time integral, we consider two cases separately: the supremum over  $q \leq N$ , and the supremum over  $q > N$ . We then use (3.2.3) to handle the supremum over  $q > N$ . We write

$$\|\omega(t)\|_{B_{p,\infty}^{\sigma(t)}} \leq \|\omega(t_0)\|_{B_{p,\infty}^{\sigma(t_0)}} + I_1 + I_2,$$

where

$$I_1 = \sup_{q \leq N} \left\{ \int_{t_0}^t C_1(e^{C\tau} + \|S_{q-1}\nabla v(\tau)\|_{L^\infty}) 2^{q\sigma(t)-q\sigma(\tau)} \|\omega(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} d\tau \right\},$$

and

$$I_2 = \sup_{q > N} \left\{ \int_{t_0}^t C_1(e^{C\tau} + \|S_{q-1}\nabla v(\tau)\|_{L^\infty}) 2^{q\sigma(t)-q\sigma(\tau)} \|\omega(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} d\tau \right\}.$$

To bound  $I_2$ , we apply (3.2.3), and we conclude that

$$I_2 \leq \int_{t_0}^t C e^{C\tau} \|\omega(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} d\tau + \sup_{q > N} \left\{ \int_{t_0}^t C_1 \epsilon q 2^{q\sigma(t)-q\sigma(\tau)} \|\omega(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} d\tau \right\}. \quad (3.4.1)$$

To handle the second integral in (3.4.1), we integrate by parts. Letting  $\sigma(t) = \text{sexp}(-\frac{2C_1}{s}\epsilon t)$  (this is the definition of  $\sigma(t)$  with  $C = 2C_1$ ), we let  $u = e^{\frac{2C_1}{s}\epsilon\tau}$  and  $dv = C_1\epsilon q e^{-\frac{2C_1}{s}\epsilon\tau} 2^{q\sigma(t)-q\sigma(\tau)} d\tau$ . Then, substituting  $u$  and  $dv$  into the second integral in (3.4.1), and recognizing that  $du$  and  $v$  are positive for all  $\tau \in [t_0, t]$ , we write

$$\begin{aligned} \sup_{q>N} \{ \sup_{\tau \in [t_0, t]} \|\omega(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} \int_{t_0}^t u dv \} &\leq \sup_{q>N} \{ \sup_{\tau \in [t_0, t]} \|\omega(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} (uv|_{t_0}^t) \} \\ &= \sup_{\tau \in [t_0, t]} \|\omega(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} e^{\frac{2C_1}{s}\epsilon t} \frac{1}{2 \ln 2}. \end{aligned} \quad (3.4.2)$$

We now bound the first time integral on the right-hand side of (3.4.1) by

$$C e^{Ct} \sup_{\tau \in [t_0, t]} \|\omega(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} (t - t_0). \quad (3.4.3)$$

Combining (3.4.3) with (3.4.2), we conclude that

$$I_2 \leq \sup_{\tau \in [t_0, t]} \|\omega(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} \{ e^{\frac{C}{s}\epsilon t} \frac{1}{2 \ln 2} + C e^{Ct} (t - t_0) \}. \quad (3.4.4)$$

To bound  $I_1$ , we first observe that  $\|S_{q-1} \nabla v(\tau)\|_{L^\infty} \leq q \|\nabla v(\tau)\|_{C_*^0}$ . Then, bounding  $\|\nabla v(\tau)\|_{C_*^0}$  by  $C(\|\omega^0\|_{L^\infty} + \|\omega^0\|_{L^{p_0}})$ , for  $p_0 \in (1, \infty)$ , and recognizing that  $2^{q(\sigma(t)-\sigma(\tau))} \leq 1$  for all  $q$ , we conclude that

$$I_1 \leq C N e^{Ct} (t - t_0) \sup_{\tau \in [t_0, t]} \|\omega(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}}. \quad (3.4.5)$$

We now combine our estimates for  $I_1$  and  $I_2$  given in (3.4.4) and (3.4.5), which gives

$$\sup_{\tau \in [t_0, t]} \|\omega(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} \leq \|\omega(t_0)\|_{B_{p,\infty}^{\sigma(t_0)}} + C^* \sup_{\tau \in [t_0, t]} \|\omega(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}},$$

where we let

$$C^* = e^{\frac{C}{s}\epsilon t} \frac{1}{2 \ln 2} + C N e^{Ct} (t - t_0). \quad (3.4.6)$$

To complete the proof, we must make the constant  $C^* < 1$ . Fix  $t > 0$ . Given this  $t$ , choose  $\epsilon > 0$  small enough to ensure that  $\frac{e^{\frac{C}{s} \epsilon t}}{2 \ln 2} < 1$ . Depending on our choice of  $t$  and  $\epsilon$ ,  $N = N(t, \epsilon)$  may be very large. Given this  $N$ , make  $t - t_0$  small enough so that  $C^* < 1$ . Note that, under these assumptions,  $C^* < 1$  when we are working on an interval of length less than or equal to  $t - t_0$ , as long as the right endpoint of the interval is less than or equal to  $t$ . We therefore break  $[0, t]$  into a finite number  $M = M(t, \epsilon)$  of intervals of length  $t - t_0$ , and we apply a bootstrapping argument. This gives

$$\sup_{\tau \in [0, t]} \|\omega(\tau)\|_{B_{p, \infty}^{\sigma(\tau)}} \leq C^M \|\omega^0\|_{B_{p, \infty}^s},$$

where  $C = \frac{1}{1 - C^*}$ , and  $C^M$  depends on  $t$  and  $\epsilon$ . More precisely, larger initial choice of  $t$  and smaller choice of  $\epsilon$  result in larger  $M$  and thus larger  $C^M$ .

This completes the proof for regularity of vorticity. To show that this implies regularity of the velocity, we need the following estimate:

**Lemma 3.4.2.** *Let  $v^0 \in B_{p, \infty}^{s+1}(\mathbb{R}^2)$ . Then there exists two positive constants  $C_0$  and  $C_1$  such that*

$$\|v(t)\|_{B_{p, \infty}^{\sigma(t)+1}} \leq C_0 e^{C_1 t} + \|\omega(t)\|_{B_{p, \infty}^{\sigma(t)}}.$$

*Proof.* We follow the proof of Lemma 6.2 in [20]. We begin by writing

$$\begin{aligned} \|v(t)\|_{B_{p, \infty}^{\sigma(t)+1}} &\leq \|\Delta_{-1} v(t)\|_{B_{p, \infty}^{\sigma(t)+1}} + \|(Id - \Delta_{-1}) \Delta_j v(t)\|_{B_{p, \infty}^{\sigma(t)+1}} \\ &\leq \|\Delta_{-1} v(t)\|_{L^p} + \|\omega(t)\|_{B_{p, \infty}^{\sigma(t)}}, \end{aligned} \tag{3.4.7}$$

where we used Lemma 2.3.8 in Section 2.3.3 on the high frequency term. We bound  $\|\Delta_{-1} v(t)\|_{L^p}$  as we did in Lemma 2.3.10. This gives the lemma.  $\square$

This completes the proof of Theorem 3.4.1.  $\square$

### 3.4.2 Besov regularity of the flow

In this section, we prove the following theorem:

**Theorem 3.4.3.** *Let  $v^0$  and  $\omega^0$  satisfy the assumptions of Theorem 3.4.1. Let  $g(t, x)$  be the measure-preserving homeomorphism satisfying  $\partial_t g(t, x) = v(t, g(t, x))$ . Define  $h(t, x) = g(t)^{-1}(x) - x$ . Then, for fixed  $\delta > 0$ ,  $\|h(t)\|_{B_{p,\infty}^{\sigma+1-\delta}}$  belongs to  $L_{loc}^\infty(\mathbb{R}^+)$ .*

*Proof.* We show that  $h(t) \in B_{p,\infty}^{\sigma'(t)}(\mathbb{R}^2)$ , where  $\sigma'(t) = \sigma(t) + 1 - \delta$ , and  $\delta$  is the Holder exponent of  $h(t)$  (see Theorem 3.1.3). The proof of Theorem 3.4.3 is similar to that for Theorem 3.4.1. However, we must deal with the extra term which shows up in the commutator estimate given in Proposition 3.3.3.

We begin by applying Proposition 3.3.3 to (3.3.17), where, once again, we drop the dependence of  $C_1$  on  $\sigma'$ , and we introduce the dependence of  $C_2$  on time, as given in Remark 3.3.1. This gives

$$\begin{aligned} \|\Delta_q h(t)\|_{L^p} &\leq \|\Delta_q h(t_0)\|_{L^p} + \int_{t_0}^t C_1 (e^{C\tau} + \|S_{q-1} \nabla v(\tau)\|_{L^\infty}) 2^{-q\sigma'(\tau)} \|h(\tau)\|_{B_{p,\infty}^{\sigma'(\tau)}} d\tau \\ &\quad + \int_{t_0}^t \left\{ \sum_{q'=q-4}^{q+4} (\|h(\tau)\|_{C^{1-\delta}} \|\Delta_{q'} v(\tau)\|_{L^p} C(\delta) 2^{q'\delta}) + \|\Delta_q v(\tau)\|_{L^p} \right\} d\tau. \end{aligned}$$

We now multiply both sides of the inequality by  $2^{q\sigma'(t)}$  and take the supremum

over  $q$  to bound  $\|h(t)\|_{B_{p,\infty}^{\sigma'(t)}}$  above by

$$\begin{aligned} & \|h(t_0)\|_{B_{p,\infty}^{\sigma'(t_0)}} + \sup_q \left\{ \int_{t_0}^t C_1 (e^{C\tau} + \|S_{q-1} \nabla v(\tau)\|_{L^\infty}) 2^{q(\sigma'(t)-\sigma'(\tau))} \|h(\tau)\|_{B_{p,\infty}^{\sigma'(\tau)}} d\tau \right\} \\ & + \int_{t_0}^t \{C(\delta) \|h(\tau)\|_{C^{1-\delta}} \|v(\tau)\|_{B_{p,\infty}^{\sigma(\tau)+1}} + \|v(\tau)\|_{B_{p,\infty}^{\sigma(\tau)+1}}\} d\tau. \end{aligned}$$

Here we used the fact that  $\sigma'(t) = \sigma(t) + 1 - \delta$ , with  $\delta > 0$ . The constant  $C(\delta)$  now depends on  $\sigma(\tau)$ , but it is uniformly bounded for all  $\sigma(\tau) \in (0, 2)$ .

We replace  $\sigma'(t) - \sigma'(\tau)$  with  $\sigma(t) - \sigma(\tau)$  in the first time integral, and we get

$$\|h(t)\|_{B_{p,\infty}^{\sigma'(t)}} \leq \|h(t_0)\|_{B_{p,\infty}^{\sigma'(t_0)}} + J_1 + J_2,$$

where

$$J_1 = \sup_q \left\{ \int_{t_0}^t C_1 (e^{C\tau} + \|S_{q-1} \nabla v(\tau)\|_{L^\infty}) 2^{q(\sigma(t)-\sigma(\tau))} \|h(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} d\tau \right\},$$

and

$$J_2 = \int_{t_0}^t \{C(\delta) \|h(\tau)\|_{C^{1-\delta}} \|v(\tau)\|_{B_{p,\infty}^{\sigma(\tau)+1}} + \|v(\tau)\|_{B_{p,\infty}^{\sigma(\tau)+1}}\} d\tau.$$

The argument for dealing with  $J_1$  is identical to the argument we used to handle  $I_1$  and  $I_2$  when proving the first part of Theorem 3.4.1. Following this approach, we conclude that

$$\sup_{\tau \in [t_0, t]} \|h(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} \leq \|h(t_0)\|_{B_{p,\infty}^{\sigma'(t_0)}} + C^* \sup_{\tau \in [t_0, t]} \|h(\tau)\|_{B_{p,\infty}^{\sigma(\tau)}} + J_2,$$

where  $C^*$  is given by (3.4.6). Arguing as we did with  $\omega$ , we make  $C^* < 1$  on a sufficiently short time interval and use a bootstrapping argument, as well as

the fact that  $h(0, x) = 0$ , to conclude that

$$\sup_{\tau \in [0, t]} \|h(\tau)\|_{B_{p, \infty}^{\sigma'(\tau)}} \leq C \int_0^t \{C(\delta) \|h(\tau)\|_{C^{1-\delta}} \|v(\tau)\|_{B_{p, \infty}^{\sigma(\tau)+1}} + \|v(\tau)\|_{B_{p, \infty}^{\sigma(\tau)+1}}\} d\tau. \quad (3.4.8)$$

We now observe that the right hand side of (3.4.8) is finite by Theorem 3.1.3 and by the first part of Theorem 3.4.1. This completes the proof of the theorem.  $\square$

## Chapter 4

### Vanishing viscosity in the plane

#### 4.1 Introduction and history

The problem of vanishing viscosity addresses whether or not a solution of the Navier-Stokes equations with initial data  $v^0$  converges in some norm to a solution of the Euler equations with the same initial data as viscosity tends to 0. This area of research is closely tied to uniqueness of solutions to the Euler equations, primarily because the methods used to prove uniqueness can be applied to show vanishing viscosity. Two of the most important uniqueness results in the plane are due to Yudovich. He proves in [23] the uniqueness of a solution  $v$  to the Euler equations with  $v^0 \in L^2(\mathbb{R}^2)$  and  $\omega^0 \in L^p(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$  for some  $p < \infty$ . He extends this uniqueness class in [24] to solutions with unbounded vorticity, but such that the  $L^p$ -norms of the vorticity grow sufficiently slowly (for example, like  $\log p$ ).

For Yudovich's uniqueness class with bounded vorticity, Chemin proves in [5] that the vanishing viscosity limit holds in the energy norm, and he establishes a rate of convergence:

**Theorem 4.1.1.** *Let  $v^0 \in L^2(\mathbb{R}^2)$ . Assume  $v_\nu$  is the unique solution to (NS) with initial data  $v^0$ , and  $v$  is the solution to (E) with initial data  $v^0$ . Also*

assume that  $\omega(v^0) = \omega^0$  belongs to  $L^\infty(\mathbb{R}^2) \cap L^2(\mathbb{R}^2)$ . Then  $v_\nu$  converges to  $v$  in  $L_{loc}^\infty(\mathbb{R}^+; L^2(\mathbb{R}^2))$  as  $\nu$  tends to zero. Specifically,

$$\|v_\nu - v\|_{L^\infty([0,T]; L^2)} \leq C(4\nu T)^{\frac{1}{2}} \exp(-C\|\omega^0\|_{L^2 \cap L^\infty} T).$$

In [8], Hmidi and Keraani show that the vanishing viscosity limit holds in the  $B_{\infty,1}^0$ -norm when  $v^0$  belongs to  $B_{\infty,1}^1(\mathbb{R}^2)$ . (Global well-posedness for the Euler equations under these assumptions was proved by Vishik in [20].) They prove the following theorem:

**Theorem 4.1.2.** *Assume  $v^0$  belongs to  $B_{\infty,1}^1(\mathbb{R}^2)$ , and  $\operatorname{div} v = 0$ . Let  $v_\nu$  be the solution to (NS) with initial data  $v^0$ , and let  $v$  be the solution to (E) with initial data  $v^0$ . Then the following inequality holds:*

$$\|v_\nu - v\|_{L^\infty([0,T]; B_{\infty,1}^0(\mathbb{R}^2))} \leq c(\nu T)^{1/2} (1 + \nu T)^{1/2} e^{e^c T},$$

where the constants depend on the  $B_{\infty,1}^0$ -norm of  $\omega^0$ .

## 4.2 Statement of the main results

We consider a uniqueness class in which initial vorticity is in a larger space than  $B_{\infty,1}^0(\mathbb{R}^2)$ . This uniqueness class was established by Vishik in [21]. As in [21], let  $\Gamma : \mathbb{R} \rightarrow [1, \infty)$  be a locally Lipschitz continuous monotonically nondecreasing function that satisfies conditions (i)-(iii) p. 771 of [21]. Condition (i) is that  $\Gamma = 1$  on the interval  $(-\infty, -1]$  and  $\lim_{\beta \rightarrow \infty} \Gamma(\beta) = \infty$ . For

the other (minor technical) conditions see [21]. Define the space

$$B_\Gamma = \{f \in \mathcal{S}'(\mathbb{R}^2) : \sum_{j=-1}^N \|\Delta_j f\|_{L^\infty} = O(\Gamma(N))\}$$

with the norm

$$\|f\|_\Gamma = \sup_{N \geq -1} \frac{1}{\Gamma(N)} \sum_{j=-1}^N \|\Delta_j f\|_{L^\infty}.$$

The following fundamental result for initial vorticities in  $B_\Gamma$  is from Theorems 7.1 and 8.1 of [21]:

**Theorem 4.2.1.** *Define  $\Gamma_1 : \mathbb{R} \rightarrow [1, \infty)$  by*

$$\Gamma_1(\beta) = \begin{cases} 1, & \beta < -1, \\ (\beta + 2)\Gamma(\beta), & \beta \geq -1 \end{cases}$$

*and add the assumption (on  $\Gamma$ ) that  $\Gamma_1$  is convex. Finally, assume that  $\Gamma$  satisfies*

$$(\beta + 2)\Gamma'(\beta) \leq C \tag{4.2.1}$$

*for almost all  $\beta \in [-1, \infty)$ . Given initial vorticity  $\omega^0$  in  $B_\Gamma \cap L^{p_0} \cap L^{p_1}$  with  $1 < p_0 < 2 < p_1 \leq \infty$  there exists a short-time solution to (E) unique in the class of vorticities lying in  $L^\infty([0, T]; L^{p_0} \cap L^{p_1}) \cap C_{w^*}([0, T]; B_{\Gamma_1})$ . With the added assumption that*

$$\Gamma'(\beta)\Gamma_1(\beta) \leq C \tag{4.2.2}$$

*for almost all  $\beta \geq -1$ , there exists a solution to (E) unique in the class of vorticities lying in  $L_{loc}^\infty([0, \infty); L^{p_0} \cap L^{p_1}) \cap C_{w^*}([0, \infty); B_{\Gamma_1})$ . Here,  $C_{w^*}$  is the space of weak\*-continuous functions (see [21] for details).*

Observe that the vorticity degrades immediately in that (as far as is known) it belongs to a larger space at all positive times than it does at time zero.

**Remark 4.2.1.** In Theorem 4.2.2, Corollary 4.2.3, and Corollary 4.2.4 below, for the case where  $\lim_{n \rightarrow \infty} \Gamma(n) = \infty$ , the symbol  $C$  represents an unspecified *absolute* constant (that is, independent of the initial data). For the case where  $\Gamma(n)$  is bounded in  $n$ , the constant  $C$  depends on both the  $L^2$ -norm and the  $B_{\infty,1}^0$ -norm of initial vorticity. This dependence arises in Equation (4.3.6) below.

We now state our main result.

**Theorem 4.2.2.** *Let  $\Gamma : \mathbb{R} \rightarrow [0, \infty)$  (without making any of the assumptions on  $\Gamma$  of [21]) and assume that  $v^0$  is in  $L^2(\mathbb{R}^2)$  with  $\omega^0 = \omega(u^0)$  in  $B_\Gamma(\mathbb{R}^2) \cap L^2(\mathbb{R}^2)$ . Then there exists a unique solution  $v_\nu$  to (NS) and a (not necessarily unique) solution  $v$  to (E), both lying in  $L^\infty([0, \infty); H^1(\mathbb{R}^2))$ . For any such  $v$ ,*

$$\|v_\nu - v\|_{L^\infty([0, T]; L^2(\mathbb{R}^2))} \leq C(\nu T)^{1/2} \|\omega^0\|_{L^2} \exp(e^{C\alpha T \Gamma(-\log(\nu T)/2)}) \quad (4.2.3)$$

for all  $T > 0$ , where  $\alpha = \|\omega^0\|_{B_\Gamma}$ .

*Proof.* The existence of a global-in-time solution to (E) with vorticity in  $L^\infty([0, \infty); L^p(\mathbb{R}^2))$  for  $\omega^0$  in  $L^p(\mathbb{R}^2)$ ,  $p > 1$ , is due to Yudovich in [23] (see, for instance, Theorem 4.1 p. 126 of [14]). The existence and uniqueness of solutions to (NS) lying in  $L^\infty([0, T]; L^2(\mathbb{R}^2)) \cap L^2([0, T]; L^2(\mathbb{R}^2))$  for  $v^0$  in  $L^2(\mathbb{R}^2)$

is classical (see, for instance, Theorems III.3.1 and III.3.2 of [17]). Because our solutions to  $(NS)$  are in the whole plane, all  $L^p$ -norms of the vorticity are non-increasing, so, in fact,  $v_\nu$  lies in  $L^\infty([0, \infty); H^1(\mathbb{R}^2))$ .

The proof of Equation (4.2.3) is contained in the sections that follow. □

It is possible to loosen the finite energy requirement in Theorem 4.2.2 that  $v^0$  lie in  $L^2(\mathbb{R}^2)$ , allowing it to lie, for instance, in the space  $E_m$  of [6].

Without restrictions on  $\Gamma$  it is of course possible that the right-hand side of Equation (4.2.3) will not go to zero with  $\nu$ . In order to establish the vanishing viscosity limit,  $\Gamma(n)$  cannot grow any faster than  $C \log n$ . We have the following immediate corollary of Theorem 4.2.2:

**Corollary 4.2.3.** *When  $\Gamma(n) = O(\log n)$ ,  $v_\nu \rightarrow v$  in  $L^\infty([0, T]; L^2(\mathbb{R}^2))$  for  $T < (C\alpha)^{-1}$ , with*

$$\begin{aligned} & \|v_\nu - v\|_{L^\infty([0, T]; L^2(\mathbb{R}^2))} \\ & \leq C \|\omega^0\|_{L^2} (\nu T)^{1/2} \exp\left(\left(-\frac{1}{2} \log(\nu T)\right)^{C\alpha T}\right). \end{aligned} \quad (4.2.4)$$

In Corollary 4.2.4, we extend the class of solutions for which both existence and uniqueness of solutions to  $(E)$  can be demonstrated globally in time. Note that we obtain uniqueness in Corollary 4.2.4 in spite of lacking knowledge of whether the solution to  $(E)$  remains in the class  $L^\infty([0, T]; B_{\Gamma_1}(\mathbb{R}^2))$  for arbitrarily large  $T$ , this being (almost) the class for which Vishik demonstrates uniqueness in [21] (see the comment on p. 771 of [21]).

**Corollary 4.2.4.** *When  $\Gamma(n) = O(\log^\kappa n)$  with  $0 \leq \kappa < 1$ , the solution  $v$  to (E) is unique in  $L^\infty([0, \infty); H^1(\mathbb{R}^2))$ . Also,  $v_\nu \rightarrow v$  in  $L_{loc}^\infty([0, \infty); L^2(\mathbb{R}^2))$ , and for all  $T > 0$ ,*

$$\|v_\nu - v\|_{L^\infty([0, T]; L^2(\mathbb{R}^2))} \leq C(\nu T)^{1/2} \|\omega^0\|_{L^2} \exp(e^{C\alpha T \log^\kappa(-\log(\nu T)/2)}). \quad (4.2.5)$$

*Proof.* The rate in Equation (4.2.5) follows immediately from Theorem 4.2.2. By Equation (4.2.5), any solution  $v$  to (E) lying in  $L^\infty([0, \infty); H^1(\mathbb{R}^2))$  is the strong limit in  $L_{loc}^\infty([0, \infty); L^2(\mathbb{R}^2))$  of the solutions  $v_\nu$  to (NS); since strong limits are unique, we conclude that the solution  $v$  is unique.  $\square$

In Corollary 4.2.4, one can show that a solution to (E) in  $L^\infty([0, \infty); H^1(\mathbb{R}^2))$  is unique without using the vanishing viscosity limit. Indeed, given a solution  $v$  to (E) with initial data  $v^0$ , we construct in the proof of Theorem 4.2.2 a sequence of  $C^\infty(\mathbb{R}^2)$  solutions  $v_n$  to (E) with initial data  $S_n v^0$ . We then show that  $\omega^0 \in B_\Gamma(\mathbb{R}^2)$  implies  $\|v_n - v\|_{L^\infty([0, T]; L^2(\mathbb{R}^2))}$  goes to 0 as  $n$  approaches infinity (see Equation (4.3.3) and Equation (4.3.7) in the sections that follow), where  $\Gamma$  satisfies the conditions in Corollary 4.2.4. Since the sequence  $v_n$  is uniquely determined by the initial data  $v^0$ , two solutions to (E) with the same initial data and initial vorticity in  $B_\Gamma(\mathbb{R}^2)$  will have the same approximating sequence and will therefore be equal on  $[0, T]$ .

The restriction Equation (4.2.1) on  $\Gamma$  ensures that  $\Gamma(N)$  grows no faster than  $C \log N$  for large  $N$ . Therefore, Corollaries 4.2.3 and 4.2.4 establish a rate of convergence for the entire short time existence and uniqueness class in

[21]. Similarly, the assumption Equation (4.2.2) on  $\Gamma$  ensures that  $\Gamma(N)$  grows no faster than  $C \log^{\frac{1}{2}} N$  for large  $N$ . Therefore, Corollary 4.2.4 establishes a rate of convergence for the entire global existence and uniqueness class in [21] as well.

### 4.2.1 A comparison of rates of convergence

Note that when  $\kappa = 0$ , our result reduces to the case where  $\omega^0 \in B_{\infty,1}^0(\mathbb{R}^2)$ , and our rate of convergence becomes

$$\|v_\nu - v\|_{L^\infty([0,T];L^2(\mathbb{R}^2))} \leq C(\nu T)^{1/2} \|\omega^0\|_{L^2} \exp(e^{C\alpha T}). \quad (4.2.6)$$

Since  $B_{\infty,1}^0(\mathbb{R}^2) \subseteq L^\infty(\mathbb{R}^2)$ , we would expect this rate to be faster than that established by Chemin in 4.1.1, which it is. Chemin's rate is substantially better than that of 4.2.4 and of 4.2.6 with  $0 < \kappa < 1$ ; however, the two spaces  $B_{O(\log^\kappa n)}(\mathbb{R}^2)$  and  $L^\infty(\mathbb{R}^2)$  are not comparable for  $0 < \kappa \leq 1$ , since the vorticity can be unbounded for the first space while  $\Gamma(n) = O(n)$  for the second.

Finally, the rate established by Hmidi and Keraani in Theorem 4.1.2 is the same as that in Equation (4.2.5) up to dependence of constants on time and initial data, although they show convergence in a different norm.

## 4.3 Proof of the main theorem

We now begin the proof of Theorem 4.2.2. Let

$$v_n = \text{the solution to } (E) \text{ with initial velocity } v_n^0,$$

where  $v_n^0$ ,  $n = 1, 2, \dots$ , is a divergence-free initial velocity smoothed to lie in  $C^\infty(\mathbb{R}^2)$  and such that  $v_n^0 \rightarrow v^0$  in  $L^2(\mathbb{R}^2)$  as  $n \rightarrow \infty$ . Letting

$$X = L^\infty([0, T]; L^2(\mathbb{R}^2))$$

we have, for any solution  $v$  to (E) in  $L^\infty([0, \infty); H^1(\mathbb{R}^2))$ ,

$$\|v_\nu - v\|_X \leq \|v_\nu - v_n\|_X + \|v - v_n\|_X.$$

To bound the first term on the right hand side, we use an energy estimate which can be found in [13]:

$$\begin{aligned} \|v_\nu(t) - v_n(t)\|_{L^2}^2 &\leq C\nu t \|\omega^0\|_{L^2} \|\omega(v_n^0)\|_{L^2} + \|v^0 - v_n^0\|_{L^2}^2 \\ &\quad + 2 \int_0^t \int_{\mathbb{R}^2} |v_\nu(s, x) - v_n(s, x)|^2 |\nabla v_n(s, x)| \, dx \, ds. \end{aligned} \quad (4.3.1)$$

As long as we insure that the initial velocity is smoothed in such a way that

$$\|\omega(v_n^0)\|_{L^2} \leq C \|\omega^0\|_{L^2} \quad (4.3.2)$$

we can conclude from Gronwall's inequality and (4.3.1) that

$$\|v_\nu(t) - v_n(t)\|_{L^2}^2 \leq (C\nu t \|\omega^0\|_{L^2}^2 + \|v^0 - v_n^0\|_{L^2}^2) e^{2 \int_0^t \|\nabla v_n\|_{L^\infty}}$$

so

$$\|v_\nu - v_n\|_X \leq ((C\nu T)^{1/2} \|\omega^0\|_{L^2} + \|v^0 - v_n^0\|_{L^2}) e^{\int_0^T \|\nabla v_n\|_{L^\infty}},$$

using  $(A^2 + B^2)^{1/2} \leq A + B$  for  $A, B \geq 0$ .

The energy argument for bounding  $\|v - v_n\|_X$  is identical except that the term involving  $\nu$  is absent and of course we have  $v$  in place of  $v_\nu$ . (In

this energy argument, although the norm of  $v(t)$  in  $H^1(\mathbb{R}^2)$  does not appear, the membership of  $v(t)$  in  $H^1(\mathbb{R}^2)$  for almost all  $t$  is required to insure the vanishing of one of the two nonlinear terms, so we *are* using the membership of  $v$  in  $L^\infty([0, \infty); H^1(\mathbb{R}^2))$ .

We thus have

$$\|v - v_n\|_X \leq \|v^0 - v_n^0\|_{L^2} e^{\int_0^T \|\nabla v_n\|_{L^\infty}} \quad (4.3.3)$$

and so

$$\begin{aligned} \|v_\nu - v\|_X &\leq (C\nu T)^{1/2} \|\omega^0\|_{L^2} e^{\int_0^T \|\nabla v_n\|_{L^\infty}} \\ &\quad + 2\|v^0 - v_n^0\|_{L^2} e^{\int_0^T \|\nabla v_n\|_{L^\infty}}. \end{aligned} \quad (4.3.4)$$

Now suppose we can show that for some sequence  $(v_n^0)_{n=1}^\infty$  of approximations to  $v^0$  satisfying Equation (4.3.2),

$$\|v^0 - v_n^0\|_{L^2} e^{\int_0^T \|\nabla v_n\|_{L^\infty}} \rightarrow 0 \text{ as } \nu \rightarrow 0. \quad (4.3.5)$$

Then letting  $n = f(\nu)$  with  $f(\nu) \rightarrow \infty$  as  $\nu \rightarrow 0$ , the second term in Equation (4.3.4) will vanish with the viscosity. By choosing  $f$  to increase to infinity sufficiently slowly, we can always make the first term in Equation (4.3.4) vanish with the viscosity as well. Thus, to establish the vanishing viscosity limit, we need only show that Equation (4.3.5) holds; to determine a bound on the rate of convergence, however, we must choose the function  $f$  explicitly.

What we have done is in effect decouple the vanishing viscosity limit from the Navier-Stokes equations and from the viscosity itself. Also, we have yet to use the information we gain from  $\omega^0$  lying in  $B_\Gamma$ ; this information

is encoded in the approximate solution  $v_n$  and will be exploited in the next section.

### 4.3.1 Convergence in Equation (4.3.5)

To smooth the initial velocity let

$$v_n^0 = S_n v^0.$$

Then  $\omega_n^0 = S_n \omega^0$  and Equation (4.3.2) is satisfied. Also,

$$\begin{aligned} \|v^0 - v_n^0\|_{L^2} &= \|(\text{Id} - S_n)v^0\|_{L^2} = \left\| \sum_{q=n+1}^{\infty} \Delta_q v^0 \right\|_{L^2} \leq \sum_{q=n+1}^{\infty} \|\Delta_q v^0\|_{L^2} \\ &\leq C \sum_{q=n+1}^{\infty} 2^{-q} \|\Delta_q \nabla v^0\|_{L^2} \\ &\leq C \left( \sum_{q=n+1}^{\infty} 2^{-2q} \right)^{1/2} \left( \sum_{q=n+1}^{\infty} \|\Delta_q \nabla v^0\|_{L^2}^2 \right)^{1/2} \\ &\leq C 2^{-n} \left( \sum_{q=n+1}^{\infty} \|\Delta_q \omega^0\|_{L^2}^2 \right)^{1/2} \leq C \|\omega^0\|_{L^2} 2^{-n}, \end{aligned}$$

where we used Minkowski's inequality, Bernstein's inequality, and the Cauchy-Schwarz inequality. From Lemma 4.3.1, below,

$$\begin{aligned} \|\nabla v_n(t)\|_{L^\infty} &\leq C \left( \|\omega_n^0\|_{L^2} + \|\omega_n^0\|_{B_{\infty,1}^0} \right) e^{Ct\|\omega_n^0\|_{B_{\infty,1}^0}} \\ &\leq C \left( \|\omega^0\|_{L^2} + \alpha\Gamma(n) \right) e^{Ct\alpha\Gamma(n)} \leq C\alpha\Gamma(n) e^{C\alpha t\Gamma(n)}, \end{aligned} \tag{4.3.6}$$

where  $\alpha = \|\omega^0\|_{B_\Gamma}$ . When  $\lim_{n \rightarrow \infty} \Gamma(n) = \infty$ , Equation (4.3.6) holds for an absolute constant  $C$  for all sufficiently large  $n$ ; it holds for all  $n$  for a constant that depends upon the initial vorticity. (See Remark (4.2.1)). This applies as

well to the inequalities that follow. Also, in Equation (4.3.6) we used

$$\|\omega_n^0\|_{B_{\infty,1}^0} = \sum_{q \geq -1} \|\Delta_q \omega_n^0\|_{L^\infty} \leq \sum_{q=-1}^{n+1} \|\Delta_q \omega^0\|_{L^\infty} \leq \alpha \Gamma(n).$$

Thus,

$$\int_0^T \|\nabla v_n(t)\|_{L^\infty} \leq \frac{C\alpha\Gamma(n)}{C\alpha\Gamma(n)} (e^{C\alpha T\Gamma(n)} - 1) \leq e^{C\alpha T\Gamma(n)}$$

and

$$\|v^0 - v_n^0\|_{L^2} e^{\int_0^T \|\nabla v_n\|_{L^\infty}} \leq C \|\omega^0\|_{L^2} 2^{-n} \exp(e^{C\alpha T\Gamma(n)}). \quad (4.3.7)$$

To bound the rate of convergence of  $v_\nu$  to  $v$ , we must decide how to choose  $n$  as a function of  $\nu$  in Equation (4.3.4). Using Equation (4.3.7), we have

$$\|v_\nu - v\|_X \leq C \|\omega^0\|_{L^2} ((\nu T)^{1/2} + 2^{-n}) \exp(e^{C\alpha T\Gamma(n)}).$$

Viewing this as a sum of two rates, when  $n = -(1/2) \log(\nu T)$  the two rates are equal. If  $n$  increases more rapidly as  $\nu \rightarrow 0$  then the first term decreases more slowly as  $\nu \rightarrow 0$ ; if  $n$  increases more slowly as  $\nu \rightarrow 0$  then the second term decreases more slowly as  $\nu \rightarrow 0$ . Since the slower decreasing of the two terms limits the convergence rate, we conclude that letting  $n = -(1/2) \log(\nu T)$  optimizes the convergence rate, giving the bound in Theorem 4.2.2 and completing its proof.

**Lemma 4.3.1.** *Let  $v$  be a  $C^\infty$ -solution to (E) with initial velocity  $v^0$ , where  $\omega^0$  is in  $L^{p_0}(\mathbb{R}^2) \cap B_{\infty,1}^0(\mathbb{R}^2)$ , with  $p_0$  in  $(1, \infty)$ . Then*

$$\|\nabla v(t)\|_{L^\infty} \leq C \left( \|\omega^0\|_{L^{p_0}} + \|\omega^0\|_{B_{\infty,1}^0} \right) e^{Ct\|\omega^0\|_{B_{\infty,1}^0}}.$$

*Proof.* We have,

$$\begin{aligned}
\|\nabla v(t)\|_{L^\infty} &\leq \|\Delta_{-1}\nabla v(t)\|_{L^\infty} + \sum_{q \geq 0} \|\Delta_q \nabla v(t)\|_{L^\infty} \\
&\leq C \|\Delta_{-1}\omega(t)\|_{L^{p_0}} + C \sum_{q \geq 0} \|\Delta_q \omega(t)\|_{L^\infty} \\
&\leq C \|\omega^0\|_{L^{p_0}} + C \|\omega(t)\|_{B_{\infty,1}^0}.
\end{aligned}$$

Here we used Bernstein's inequality with the Calderon-Zygmund inequality for the first term and Lemma 2.3.8 for the sum.

From Theorem 4.2 of [20],

$$\|\omega(t)\|_{B_{\infty,1}^0} \leq C(1 + \log(\|g(t)\|_{lip}\|g^{-1}(t)\|_{lip}))\|\omega^0\|_{B_{\infty,1}^0},$$

where  $g$  is the flow associated to  $v$ ; that is,

$$g(t, x) = x + \int_0^t v(s, g(s, x)) ds.$$

It follows from Gronwall's inequality that

$$\|g(t)\|_{lip}, \|g^{-1}(t)\|_{lip} \leq \exp \int_0^t \|\nabla v(s)\|_{L^\infty} ds.$$

Combining the three inequalities above gives

$$\|\nabla v(t)\|_{L^\infty} \leq C\|\omega^0\|_{L^{p_0}} + C \left(1 + 2 \int_0^t \|\nabla v(s)\|_{L^\infty} ds\right) \|\omega^0\|_{B_{\infty,1}^0},$$

and the proof is completed by another application of Gronwall's inequality.  $\square$

In the proof of Lemma 4.3.1 we used the existence of a flow associated with a smooth solution to  $(E)$ , which allowed us to apply Theorem 4.2 of

[20]. This is where our approach differs markedly from that of Vishik's in [21], where required properties of the flow are inferred from the membership of the vorticity in the spaces  $L^{p_0}(\mathbb{R}^2) \cap B_\Gamma(\mathbb{R}^2)$  and  $L^{p_0}(\mathbb{R}^2) \cap L^{p_1}(\mathbb{R}^2)$  and where the constraints on the values of  $p_0$  and  $p_1$  of Theorem 4.2.1 are required. Vishik also requires that  $p_0 < 2$  so that the velocity can be recovered uniquely from the vorticity using the Biot-Savart law, since he uses the vorticity formulation of a weak solution to  $(E)$  in [21]. By contrast, in Theorem 4.2.2 we in effect require that  $p_0 = p_1 = 2$ , so that we can make the basic energy argument in Section 4.3.

It is also possible to prove Lemma 4.3.1 using an argument like that in [8].

## Bibliography

- [1] H. Bahouri and J.-Y. Chemin. Equations de transport relatives à des champs de vecteurs non-lipschitziens et mécanique des fluides. *Arch Rational Mech. Anal*, 127:159-181, 1994.
- [2] A. Bertozzi and A. Majda. *Vorticity and incompressible flow*. Cambridge University Press, Cambridge, 2002.
- [3] J.-M. Bony. Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires. *Annales de l'Ecole Normale Supérieure*, 14:209-246, 1981.
- [4] D. Chae. On the Euler equations in the critical Triebel-Lizorkin spaces. *Arch. Rational Mech. Anal.*, 170: 185-210, 2003.
- [5] J.-Y. Chemin. A remark on the inviscid limit for two-dimensional incompressible fluids. *Comm. Partial Differential Equations*, 21(11-12):1771–1779, 1996.
- [6] J.-Y. Chemin. *Perfect incompressible fluids*, volume 14 of *Oxford Lecture Series in Mathematics and its Applications*. The Clarendon Press Oxford University Press, New York, 1998. Translated from the 1995 French original by Isabelle Gallagher and Dragos Iftimie.

- [7] M. Frazier, B. Jawerth, and G. Weiss. *Littlewood-Paley theory and the study of function spaces*, *Regional Conference Series in Mathematics*, American Mathematical Society, Providence 1991.
- [8] T. Hmidi and S. Keraani. Incompressible viscous flows in borderline Besov spaces. *Preprint de l'IRMAR, Université Rennes1*.
- [9] B. Jawerth. Some observations on Besov and Triebel-Lizorkin spaces. *Math. Scand.*, 40: 94-104, 1977.
- [10] F. John and L. Nirenberg. On functions of bounded mean oscillation. *Comm. Pure Appl. Math.*, 14: 415-426, 1961.
- [11] T. Kato, G. Ponce. Well-posedness of the Euler and Navier-Stokes equations in the Lebesgue spaces  $L^p_s(\mathbb{R}^2)$ . *Rev. Mat. Iberoamericana*, 2:73-88, 1986.
- [12] T. Kato, G. Ponce. On nonstationary flows of viscous and ideal fluids in  $L^p_s(\mathbb{R}^2)$ . *Duke Math J.*, 55:487-499, 1987.
- [13] J. Kelliher. The inviscid limit for two-dimensional incompressible fluids with unbounded vorticity. *Math. Res. Lett.*, 11(4):519–528, 2004.
- [14] P.-L. Lions. *Mathematical topics in fluid mechanics. Vol. 1*, volume 3 of *Oxford Lecture Series in Mathematics and its Applications*. The Clarendon Press Oxford University Press, New York, 1996.

- [15] D. Sarason. Functions of vanishing mean oscillation. *Trans. Amer. Math. Soc.*, 207: 391-405, 1975.
- [16] E. Stein. *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*. Princeton University Press, Princeton, New Jersey, 1993.
- [17] R. Temam. *Navier-Stokes equations*. AMS Chelsea Publishing, Providence, RI, 2001. Theory and numerical analysis, Reprint of the 1984 edition.
- [18] A. Torchinsky. *Real-variable methods in harmonic analysis*. Academic Press, Inc, Orlando, Florida, 1986.
- [19] H. Triebel. *Theory of function spaces II*, volume 84 of *Monographs in mathematics*. Birkhäuser, Basel, 1992.
- [20] M. Vishik. Hydrodynamics in Besov spaces. *Arch. Ration. Mech. Anal.*, 145(3):197–214, 1998.
- [21] M. Vishik. Incompressible flows of an ideal fluid with vorticity in borderline spaces of Besov type. *Ann. Sci. École Norm. Sup. (4)*, 32(6):769–812, 1999.
- [22] M. Yamazaki. A quasi-homogeneous version of paradifferential operators, I. Boundedness on spaces of Besov type. *J. Fac. Sci. Univ. Tokyo Sect. I A Math.*, 33: 131-174, 1986.

- [23] V. I. Yudovich. Non-stationary flows of an ideal incompressible fluid. *Ž. Vyčisl. Mat. i Mat. Fiz.*, 3:1032–1066 (Russian), 1963.
- [24] V. I. Yudovich. Uniqueness theorem for the basic nonstationary problem in the dynamics of an ideal incompressible fluid. *Math. Res. Lett.*, 2(1):27–38, 1995.

# Vita

Elaine Marie Cozzi was born on August 2, 1978 in Falls Church, Virginia, the daughter of Katherine Marie Cozzi and Howard Allen Cozzi. She graduated from Bishop O'Connell High School in Arlington, Virginia in 1996 and received a Bachelor of Arts degree in Mathematics and Economics from the University of Virginia in May, 2000. She began her graduate studies at the University of Texas at Austin in August, 2001.

Permanent address: 301 E. 33rd St. Apt. 2  
Austin, Texas 78705

This dissertation was typeset with  $\text{\LaTeX}^\dagger$  by the author.

---

<sup>†</sup> $\text{\LaTeX}$  is a document preparation system developed by Leslie Lamport as a special version of Donald Knuth's  $\text{\TeX}$  Program.