

## Catalogue of American Amphibians and Reptiles.

SALTHE, STANLEY N. 1973. *Amphiuma means*.*Amphiuma means* Garden  
Two-toed congo eel

*Amphiuma means* Garden, 1821, p. 599. In Smith, Correspond. of Linn. Type locality not stated, but from context either Charleston, South Carolina or eastern Florida; restricted to Charleston, Charleston County, South Carolina by Schmidt, 1953:27. Holotype possibly rediscovered by Lönnberg, 1896:36, #15 labelled *Siren lacertina* in the Zoological Museum of the University of Upsala (not seen by author).

*Sireni simile* Linnaeus, 1821, p. 599. In Smith, Correspond. of Linn. Refers to an apparently lost letter from Linnaeus to Garden.

*Chrysodonta larvaformis* Mitchill, 1822:503. Type locality Savannah, Chatham County, Georgia.

*Amphiuma didactylum* Cuvier, 1827:4. Substitute name.

*Amphiuma tridactylum*: Tschudi, 1838:97 (in part).

*Sirenoideis didactyla*: Fitzinger, 1843:34. New combination.

*Amphiuma means*: Gray, 1850:55. New combination.

*Amphiuma means means*: Goin, 1938:128. See Remarks under *A. tridactylum*.

• CONTENT. No subspecies are currently recognized; see Remarks under *A. tridactylum*.

• DEFINITION. This species tends to be unicolored, with the venter only slightly lighter than the dorsum, the change in shade between the two being relatively gradual. The dark patch on the throat is not evident against the relatively dark ventral coloration. No more than two toes are present on any limb and not less than two on most. Forelimb length in body length, 44 (Baker, 1947), 47–50 (data of Hill, 1954); hindlimb length in body length, 35 (Baker, 1947), 31–34 (data of Hill, 1954). The tail base in section is circular, not compressed.

• DESCRIPTIONS. See *Amphiuma* for many references on general anatomy. Good general descriptions are given by Mitchill (1822), Harlan (1823), Cuvier (1827), Bishop (1943), and Baker (1945). The limb and girdle skeletons are described by Parker (1868), Rabl (1901), and Stoudemayer (1949), while limb development is described by Van Pée (1903, 1904), Davison (1895), Low (1929), and Hilton (1947). The male cloaca is described by Davison (1895). Female cloacal structures and spermatophores have not been described. The ova have not been described. Rough descriptions of the eggs can be found in Brimley (1944) and Weber (1944), and the egg capsules are described by Salthe (1963). Developmental stages have not been described. The hatchling (55 mm) is described by Weber (1944). Noble and Brady (1933) comment on the hatching glands. The size at transformation is given as just under 3 inches by Harlan (1825).

• ILLUSTRATIONS. Excellent paintings of adults may be found in Wagler (1833:plate 19) and Holbrook (1842:plate 30) while Bishop (1943:52) provides photographs. Baker (1947:9) gives a photograph of the ventral view of the anterior part of the body. Stoudemayer (1949:7) presents line drawings of the appendicular skeleton. Spermatophores, eggs, and developmental stages have not been figured, but Salthe (1963:163) gives a figure of the egg and its capsules. Van Pée (1903:38–39) figures limb development in a series from 40 mm to 127 mm and Low (1929:466–467) gives a figure of the same at 60 mm. A photograph of a female at the nest site with a clutch of eggs is in Neill (1971:70).


• DISTRIBUTION. The range includes the Atlantic Coastal Plain from easternmost Virginia to the Florida Everglades and continues west in the Gulf Coastal Plain from Florida to the region around New Orleans, east of the Mississippi Valley. Viosca (1923, 1926) suggested that clay soils of alluvial or loess origin in the Mississippi Valley may impose the western limit to the species range, and Baker (1947) emphasized that pH may be a major factor, with alkaline conditions being restrictive. The species is mainly a lowland form, always found below the fall line, although it may occasionally get

into low sandy pine hills. It is found in swamps, bayous, along the margins of muddy sloughs, in cypress heads and drainage ditches, in sluggish streams and wet meadows, and in muddy lakes—all waters tending toward acid pH. The presence of crayfish burrows, in which the animals hide and hunt, is an important feature of most habitats. They often burrow deep in the soil (Winslow, 1899; Knepton, 1954), which cannot, therefore, be too dense.

• FOSSIL RECORD. Brattstrom (1953) records this species from three Pleistocene localities in Florida (Pinellas, St. Lucie, and Walkulla Counties). Neill (1957) contested these records, arguing that rapid mineralization occurs in the region, and suggested that none of the records need be more than 2500 years old. Hirschfeld (1969) noted that within a given sink hole *A. means* vertebrae could be dated from any time between late Tertiary or Pleistocene age to 1964. Weigel (1962) reports this species as very common in a bed in Indian River County, Florida that ranges in age from 30,000 years (Sangamonian) to 3500 years ago. He noted that at Redick, much further north, beds of Kansan or even Illinoian age, while containing *Siren* and *Pseudobranchius*, disclosed no *Amphiuma*. Holman (1965) and Slaughter and McClure (1965) report this species from the mid-Pleistocene (Sangamonian) of eastern Texas near Houston, outside its present range. It is not clear to the present author whether or not any of these fossils (consisting of skull fragments and vertebrae) can actually be assigned to any living species.

• PERTINENT LITERATURE. Courtship and mating occur in January in Georgia (Knepton, 1954). In northern Florida and North Carolina eggs can be found in June and July (Brimley, 1910; Davison, 1895), while in southern Florida eggs in early developmental stages have been found in February, and may have been deposited in January (Weber, 1944). The latter author suggested a 5 month incubation period. The female remains with the eggs through most of the period of development coiled under or around them (Brimley, 1910; Weber, 1944). The following nest sites have been described: under large rocks in damp places (Davison, 1895); under logs in the partly dried mud of dried-up pools (Brimley, 1910); in muck at the edge of a dried up pool under a board (Weber, 1944). The snakes *Farancia abacura* and *F. erythrogrammus* are the major predators of this species (Brimley, 1944; Funderberg, 1955; Schwartz, 1957). Other than loose descriptions of habitats (see Distribution) little further seems to be known of the ecology of this animal. See under *Amphiuma* for literature on cytology, cell physiology, and physiology.

• REMARKS. See under *A. tridactylum* a discussion of the possibility of that species being a subspecies of this one.



MAP. Solid symbol indicates type-locality, open symbols mark other localities. Stars locate Pleistocene sites for this species, or *Amphiuma tridactylum* or a common ancestor (see text). Shaded area estimates total range.

• **ETYMOLOGY.** Garden did not explain the species epithet, and it seems there has been no attempt made to understand what he had in mind. Black and Dellinger (1938) apparently assumed that it referred to a surname and accordingly used the vernacular "Means' congo cel." Assuming that Garden was a mediocre Greek scholar (see Etymology under *Amphiuma*), there are several possibilities such as *meion*, to have too little, to be scant of a thing—referring to the extraordinarily small limbs. There is also *menos*, fierceness, or *mania*, frenzy—possibly in reference to the disposition of the beast when one tries to catch it. There are still other possibilities, some referring to small size (limbs) and some to large size (body).

## LITERATURE CITED

- Baker, C. L. 1945. The natural history and morphology of Amphiumae. Rep. Reelfoot Lake Biol. Sta. 9:55-91.  
 — 1947. The species of Amphiumae. J. Tennessee Acad. Sci. 22:9-21.
- Bishop, S. C. 1943. Handbook of salamanders. Comstock Publ. Co., Ithaca, 555 p.
- Black, J. D. and S. C. Dellinger. 1938. Herpetology of Arkansas, Pt. 2, the amphibians. Occ. Pap. Univ. Arkansas Mus. (2):1-30.
- Brattstrom, B. H. 1953. Records of Pleistocene reptiles and amphibians from Florida. Quart. J. Florida Acad. Sci. 16:243-248.
- Brimley, C. S. 1910. Records of some reptiles and batrachians from the southeastern United States. Proc. Biol. Soc. Washington 23:9-18.  
 — 1944. The amphibians and reptiles of North Carolina. Carolina Biol. Sup. Co., Elon College, North Carolina, 63 p.
- Cuvier, G. J. L. F. 1827. Sur le genre de reptiles batraciens, nommé *Amphiuma*, et sur une nouvelle espece de ce genre (*Amphiuma tridactylum*). Mem. Mus. Hist. Nat. Paris 14:1-14.
- Davison, A. 1895. A contribution to the anatomy and phylogeny of *Amphiuma means*. J. Morphol. 11:375-410.
- Fitzinger, L. 1843. Systema reptilium. I. Ambyglossae. Vienna, 106 p.
- Funderberg, J. B., Jr. 1955. The amphibians of New Hanover County, North Carolina. J. Elisha Mitchell Sci. Soc. 71:19-28.
- Goin, C. J. 1938. The status of *Amphiuma tridactylum* Cuvier. Herpetologica 1:127-130.
- Gray, J. E. 1850. Catalogue of the specimens of Amphibia in the collection of the British Museum. Pt. II. Batrachia Gradientia, etc. London, 72 p.
- Harlan, R. 1823. Dissection of a batrachian animal in a living state. J. Acad. Nat. Sci. Philadelphia 3:54-59.  
 — 1825. Further observations on the *Amphiuma means*. Ann. Lyceum Nat. Hist. N. Y. 1:269-270.
- Hill, I. R. 1954. The taxonomic status of the mid-Gulf Coast *Amphiuma*. Tulane Stud. Zool. 1:191-215.
- Hilton, W. A. 1947. The skeleton of *Amphiuma*. J. Ent. Zool. 39:14-16.
- Hirschfeld, S. E. 1969. Vertebrate fauna of Nichol's Hammock, a natural trap. Quart. J. Florida Acad. Sci. 31:177-189.
- Holbrook, J. E. 1842. North American herpetology. J. Dobson & Co., Philadelphia Vol. 5, 118 p.
- Holman, J. A. 1965. A small Pleistocene herpetofauna from Houston, Texas. Texas J. Sci. 17:418-423.
- Knepton, J. C., Jr. 1954. A note on the burrowing habits of the salamander *Amphiuma means means*. Copeia 1954:68.
- Lönnberg, E. 1896. Linnaean type specimens of birds, reptiles, batrachians, and fishes in the Zoological Museum of the University of Upsala. Svensk. Vet.-akad. Handl. 22:1-45.
- Low, J. W. 1929. Contributions to the development of the pelvic girdle and its related muscles in the batrachian *Amphiuma means* of Garden. Proc. Zool. Soc. London 1929:463-468.
- Mitchill, S. L. 1822. Description of a batrachian animal from Georgia, different from the reptiles of that order hitherto known. Amer. Med. Recorder 5:499-503.
- Neill, W. T. 1957. The rapid mineralization of organic remains in Florida and its bearing on supposed Pleistocene records. Quart. J. Florida Acad. Sci. 20:1-13.  
 — 1971. The last of the ruling reptiles: alligators, crocodiles, and their kin. Columbia Univ. Press, N. Y. 486 p.
- Noble, G. K. and M. K. Brady. 1933. Observations on the life history of the marbled salamander, *Ambystoma opacum* Gravenorst. Zoologica (N. Y.) 11:89-132.
- Parker, W. K. 1868. A monograph on the structure and development of the shoulder girdle and sternum in the vertebrates. Ray Society, London, 237 p.
- Rabl, G. 1901. Gedanken und Studien über Ursprung der Extremitäten. Z. Wiss. Zool. 70:474-558.
- Salthe, S. N. 1963. The egg capsules in the Amphibia. J. Morph. 113:161-171.
- Schmidt, K. P. 1953. A checklist of North American amphibians and reptiles. Amer. Soc. Ichthyol. Herpetol., 280 p.
- Schwartz, D. 1957. "Albinism" in the salamander *Amphiuma means*. Herpetologica 13:75-76.
- Slaughter, B. H. and W. L. McClure. 1965. The Sims Bayou local fauna: Pleistocene of Houston, Texas. Texas J. Sci. 17:404-417.
- Smith, J. E. 1821. A selection of the correspondence of Linnaeus and other naturalists. Longman, Hurst, Rees, Orme, & Brown, London, Vol. I, 630 p.
- Stoudemayer, M. B. 1949. The appendicular skeleton of the Amphiumae. J. Tennessee Acad. Sci. 24:4-11.
- Tschudi, J. J. 1838. Classification der Batrachier, mit Berücksichtigung der fossilen Thiere dieser Abteilung der Reptilien. Petitpierre, Neuchatel, 102 p.
- Van Pée, P. 1903. Recherches sur le développement des extrémités chez *Amphiuma* et *Necturus*. C. R. Assoc. Anat. 5:37-43.  
 — 1904. Les membres chez *Amphiuma*. Anat. Anz. 24:476-482.
- Viosca, P., Jr. 1923. An ecological study of the cold-blooded vertebrates of southeastern Louisiana. Copeia 1923:35-44.  
 — 1926. Distributional problems of the cold-blooded vertebrates of the Gulf Coastal Plain. Ecology 7:307-314.
- Wagler, J. 1833. Descriptiones et icones amphibiorum. München, Stuttgart, u. Tübingen. 36 plates.
- Weber, J. A. 1944. Observations on the life history of *Amphiuma means*. Copeia 1944:61-62.
- Weigel, R. D. 1962. Fossil vertebrates of Vero, Florida. Florida Geol. Surv. Sp. Pub. (10):1-59.
- Winslow, G. M. 1899. On the occurrence of *Amphiuma means* in Virginia. Proc. U. S. Nat. Mus. 21:379-380.
- S. N. SALTHER, BROOKLYN COLLEGE, BROOKLYN, NEW YORK 11210

Primary editor for this account, James D. Anderson.

Published 25 October 1973 by the SOCIETY FOR THE STUDY OF AMPHIBIANS AND REPTILES.