

Catalogue of American Amphibians and Reptiles.

Nowacki, A.M. and T.M. Doan. 2012. *Phyllobates terribilis*.

***Phyllobates terribilis* Myers, Daly, and Malkin 1978**

Golden Poison Dart Frog, Terrible Poison Dart Frog

Phyllobates terribilis Myers, Daly, and Malkin 1978: 313. Type-locality, "lowland rain forest at Quebrada Guanguí, about 0.5 km above its junction with Río Patia, 100–200m elevation, in upper Río Saija drainage, Department of Cauca, Colombia." Holotype, American Museum of Natural History (AMNH) 88876 (field no. CWM 11920), an adult male, collected 18–19 February 1973 by C.W. Myers and J.W. Daly (not examined by authors).

Dendrobates terribilis: Laschat, Narjes, and Overman 1994:348. Invalid emendation.

- **CONTENT.** No subspecies are recognized.
- **DEFINITION.** *Phyllobates terribilis* is a relatively large dendrobatid with males reaching up to 45 mm SVL. Females are slightly larger and can attain a maximum SVL of 47 mm. Males mature at roughly 37 mm and females at 40–41 mm SVL. The males have a shallow subgular vocal sac which is typically indicated by small, grey expansion wrinkles at the base of the throat. Males also have well-developed vocal slits on the floor of the mouth. Measured from the base, the third finger is longest and the first finger is next, followed by the roughly equal length second and fourth. The skin is smooth to finely rugose or finely granular, turning conspicuously rugose to very coarsely granular on the upper surfaces of the hind limbs. Body coloration is nearly uniform and constant despite light or temperature changes. The range of colors observed at the type-locality include pale metallic green, pale greenish yellow, pale yellow, golden yellow, golden orange, and orange (Myers et al. 1978). Another site also has a turquoise cream morph (Lötters et al. 1997). Dorsally the color is uniform along the body and limbs with few exceptions, including black digit tips, usually black edging on the lower rim of the tympanum and often black edging along the mouth and creases of limb segments. Ventrally the color is the same or slightly lighter than the dorsal color aside from the black palms, soles, and seat patch. The call of *Phyllobates terribilis* is described as a long melodious trill. Laboratory recordings show the call is comprised of a uniform train of notes produced at a rate of 13 per second. The dominant frequency is about 1800 Hz (Myers et al. 1978).

Freshly hatched larvae have an average total length of 11.1 mm and have uniformly grey bodies and throats, turning pale grey on the tail. At around stage 37, when the hind limbs have become fairly well developed, the tadpole is an average total length of 35.4 mm. At this stage the tadpole becomes much darker




FIGURE 1. Two adult *Phyllobates terribilis* in captivity (photograph by Marc van Doorn).

blackish-gray with bronze dorsal flecking. The flecking is particularly concentrated into two dorsolateral stripes that diverge from the snout, pass over the eyes, and extend to the tail base. Metamorphosis occurs at roughly 13 mm SVL and juveniles retain the dorsolateral stripes until they reach an SVL of 20 mm (Myers et al. 1978).

- **DIAGNOSIS.** *Phyllobates terribilis* can be distinguished from its sister species *Phyllobates bicolor* primarily by size and color pattern. *Phyllobates bicolor* is significantly smaller; the mean SVL for *P. terribilis* is 41.05 mm for males, 43.23 mm for females, whereas the respective values for *P. bicolor* are 36.17 mm for males and 38.9 mm for females. In *P. bicolor* the venter and extremities are darker (usually black) than the dorsum while *P. terribilis* is unicolor. *Phyllobates terribilis* also possesses higher amounts of the skin toxin batrachotoxin (Lötters et al. 1997). Other characters that can be used to distinguish the 2 species include tibia length (shorter in *P. terribilis*), head width (narrower in *P. terribilis*), and finger discs (slightly smaller in *P. terribilis*). These characters have some overlap, however, and are less reliable (Myers et al. 1978).

- **DESCRIPTIONS.** A brief description of the coloration of the frog and its use by the Chocó Indians can be found in Myers and Daly (1983). Lötters et al. (1997) included a more detailed description of the coloration and size of *Phyllobates terribilis*. Daly et al. (1980) described the levels of batrachotoxin in the skin secretions after various lengths of time. Myers et al. (1978) included thorough descriptions of external morphology, coloration, osteology, tadpoles, juveniles, and vocalizations.

- **ILLUSTRATIONS.** Photographs of three color morphs appeared in Lötters et al. (1997); other photographs are in Myers et al. (1978) and Zimmermann and Zimmermann (1985). Myers and Daly (1983) included a drawing and Zimmermann and Zimmer-



MAP. Distribution of *Phyllobates terribilis* in Colombia. The circle indicates the type-locality and the dot indicates the only other known locality.

mann (1985) included illustrations of courtship and oviposition. Zimmermann and Zimmermann (1985) also included photographs of an adult carrying a tadpole and a clutch of eggs/tadpoles. Myers et al. (1978) included photographs of a stage 40 tadpole, juveniles showing various stages of color change, and the poisoning of a dart on a frog's back. Zimmermann and Zimmermann (1985, 1992) provided sonograms of advertisement and courtship calls.

Color photographs of adults are in Andriantsiferana et al. (2005), Barceloux (2008), Bartlett (2003), Clark (2009), Killeen (2007), Moffett (1995), Obst et al. (1988), and Steven and Overman (2007), and black-and-white photographs are in Barceloux (2008), Daly

(1998), Duellman and Trueb (1986), and Myers et al. (1978).

• **DISTRIBUTION.** The known range of *Phyllobates terribilis* is very limited. It can be found in a small area in Pacific coastal Colombia, at the western foot of a northerly inclined spur of the Cordillera Occidental at an elevation of 50–200 m. The type-locality is a small tributary to the Río Saija, and other sites are in the Río Saija drainage basin or the proximity of its mouth.

• **FOSSIL RECORD.** No fossils are known.

• **PERTINENT LITERATURE.** Myers et al. (1978) provided a very comprehensive account including morphology, osteology, habitat, distribution, natural history, skin secretions, skin texture, and a description of the tadpole. Other references listed by topic include: **behavior** (Hödl and Amezcuita 2001), **captive husbandry** (Martel et al. 2011; Nogge 2004; Zimmermann and Zimmermann 1992), **conservation** (Bolívar and Lötters 2004; Caporale 1995; Furrer and Corredor 2008; Killeen 2007; Nijman and Shepherd 2010; Rueda-Almonacid 1999; Zimmermann and Zimmermann 1992, 1994), **disease** (Forzán et al. 2008; Miller et al. 2008), **distribution** (Acosta-Galvis 2000; Duellman 1999; Lötters et al. 1997; McCain and Sanders. 2010; Ruiz-Carranza et al. 1996), **morphology** (Glaw and Vences 1997; Lötters et al. 1997; Myers 1982; Myers and Daly 1983; Neuwirth et al. 1979), **reproduction** (Brown et al. 2010; Zimmermann and Zimmermann 1985), **systematics and phylogeny** (Darst et al. 2005; Glaw and Vences 1997; Grant et al. 2006; Hagman and Forsman 2003; Han 2008; Lahanas 1992; Lötters et al. 1997; Maxson and Myers 1985; Santos 2009; Santos et al. 2003; Toledo et al. 2007; Widmer et al. 2000), **toxins** (Barceloux 2008; Clarke 1997; Daly 1995, 1998a,b; Daly et al. 1980, 1987, 2003; Dumbacher et al. 2004; Gusovsky and Daly 1988; Gusovsky et al. 1986, 1987, 1988, 1992; Hollingsworth et al. 1986; Jones et al. 1999; Laschat et al. 1994; Lovenberg and Daly 1986; Müller 1996; Myers and Daly 1983; Riston




Figure 2. Wave form and audiospectrogram of *Phyllobates terribilis*. The call rate is 3.5 calls/minute; call duration is about 5.7 s; each call is comprised of notes and/or pulses ranging from 0.002–0.011 s in duration. Dominant frequency is 1361–2320 Hz. Provided by W. Ronald Heyer, Smithsonian Institution, and based on recordings by Charles W. Myers and John W. Daly provided by the American Museum of Natural History.

2006; Steven and Overman 2007; Tokuyama and Daly 1983), and **vocalizations** (Erdtmann and Amézquita 2009; Zimmermann and Zimmermann 1985, 1992).

• **REMARKS.** These frogs produce a massive quantity of the potent neurotoxin batrachotoxin; a single frog may yield up to 1900 μg (Daly et al. 1980). They are therefore at least 20 times more toxic than other *Phyllobates*, and presumably as a result bolder and less secretive (Myers and Daly 1983). The putative source for this toxin are melyrid beetles of the genus *Choresine* (Dumbacher et al. 2004). Native tribes recognized the potency of the poison and poison their darts by rubbing them along the back of a living frog, coating the dart tip (Myers and Daly 1983; Myers et al. 1978). The toxin also means that wild specimens should be handled with great care. The frogs themselves are not sensitive to the effects of batrachotoxin (Daly et al. 1980) but they are able to contract infections of *Batrachochytrium dendrobatidis*, rana-virus, and *Aeromonas* bacteria (Miller et al. 2008).

• **ETYMOLOGY.** The specific name *terribilis* is a Latin adjective meaning “terrible” or “frightful”, in reference to the potency and amount of the batrachotoxin secreted by this species. The name also alludes to the fear once evoked by the poisoned blowgun darts of a warlike people (Myers et al. 1978).

• **ACKNOWLEDGMENTS.** We thank Marc van Doorn for the use of the photograph of *P. terribilis*.

LITERATURE CITED

- Acosta-Galvis, A.R. 2000. Ranas, Salamandras y Caecilias (Tetrapoda: Amphibia) de Colombia. *Biota Colombiana* 1:289–319.
- Andriantsiferana, M., N.R. Andriamharavo, C.R. Razafindrabe, C. Harisoa, P. Rasendra, M. Garraffo, T. Spande, and J. Daly. 2005. New lipophilic alkaloids from *Mantella* frogs collected in Madagascar, p. 169–186. *Proc. 11th NAPRECA Symp. Antananarivo, Madagascar*.
- Barceloux, D.G. 2008. *Medical Toxicology of Natural Substances: Foods, Fungi, Medicinal Herbs, Plants, and Venomous Animals*. John Wiley and Sons, Inc., Hoboken, New Jersey.
- Bartlett, R.D. 2003. *Poison Dart Frogs. Facts & Advice on Care and Breeding*. Barron's Educ. Ser., Hauppauge, New York.
- Bolívar, W. and S. Lötters. 2004. *Phyllobates terribilis*. In IUCN 2011. *IUCN Red List of Threatened Species*. Version 2011.2. <www.iucnredlist.org>.
- Brown, J.L., V. Morales, and K. Summers. 2010. A key ecological trait drove the evolution of biparental care and monogamy in an amphibian. *Am. Nat.* 175: 436–446.
- Caporale, L.H. 1995. Chemical ecology: a view from the pharmaceutical industry. *Proc. Natl. Acad. Sci. USA* 92:75–82.
- Clark, V.C. 2009. Collecting arthropod and amphibian secretions for chemical analyses, p. 1–46. In W. Zhang and H. Liu (eds.), *Behavioral and Chemical Ecology*. Nova Sci. Publ., Inc.
- Clarke, B.T. 1997. The natural history of amphibian skin secretions, their normal functioning and potential medical applications. *Biol. Rev.* 72:365–379.
- Daly J.W. 1995. Alkaloids from frog skins: selective probes for ion channels and nicotinic receptors. *Braz. J. Med. Biol. Res.* 28:1033–1042.
- . 1998a. Thirty years of discovering arthropod alkaloids in amphibian skin. *J. Nat. Prod.* 61:162–172.
- . 1998b. The nature and origin of amphibian alkaloids. *Alkaloids: Chem. Biol.* 50:141–169.
- , H.M. Garraffo, T.F. Spande, V.C. Clark, J. Ma, H. Ziffer, and J.F. Cover, Jr. 2003. Evidence for an enantioselective pumiliotoxin 7-hydroxylase in dendrobatid poison frogs of the genus *Dendrobates*. *Proc. Natl. Acad. Sci. USA* 100:11092–11097.
- , C.W. Myers, J.E. Warnick, and E.X. Albuquerque. 1980. Levels of batrachotoxin and lack of sensitivity to its action in poison-dart frogs (*Phyllobates*). *Science* 208:1383–1385.
- , –, and N. Whittaker. 1987. Further classification of skin alkaloids from neotropical poison frogs (Dendrobatidae), with a general survey of toxic/noxious substances in the Amphibia. *Toxicon* 25: 1023–1095.
- Darst, C.R., P.A. Menéndez-Guerrero, L.A. Coloma, and D.C. Cannatella. 2005. Evolution of dietary specialization and chemical defense in poison frogs (Dendrobatidae): a comparative analysis. *Am. Nat.* 165:56–69.
- Duellman, W.E. 1999. Distribution patterns of amphibians in South America, p. 255–328. In W.E. Duellman (ed.), *Patterns of Distribution of Amphibians: A Global Perspective*. Johns Hopkins Univ. Press, Baltimore.
- and L. Trueb. 1986. *Amphibian Biology*. McGraw-Hill, New York.
- Dumbacher, J.P., A. Wako, S.R. Derrickson, A. Samuelson, T.F. Spande, and J.W. Daly. 2004. Melyrid beetles (*Choresine*): a putative source for the batrachotoxin alkaloids found in poison-dart frogs and toxic passerine birds. *Proc. Natl. Acad. Sci. USA* 101:15857–15860.
- Erdtmann, L. and A. Amézquita. 2009. Differential evolution of advertisement call traits in Dart-poison frogs (Anura: Dendrobatidae). *Ethology* 115: 801–811.
- Forzán, M.J., H. Gunn, and P. Scott. 2008. Chytridiomycosis in an aquarium collection of frogs: diagnosis, treatment, and control. *J. Zoo Wildl. Med.* 39:406–411.
- Furrer, S.C. and G. Corredor. 2008. Conservation of threatened amphibians in Valle del Cauca, Colombia: a cooperative project between Cali Zoological Foundation, Colombia, and Zoo Zürich, Switzerland. *Intl. Zoo Yrbk.* 42:158–164.
- Glaw, F. and M. Vences. 1997. Anuran eye colouration: definitions, variation, taxonomic implications and possible functions. *Herpetol. Bonn.* 1997: 125–138.

- Grant, T., D.R. Frost, J.P. Caldwell, R. Gagliardo, C.F.B. Haddad, P.J.R. Kok, D.B. Means, B.P. Noonan, W.E. Schargel, and W.C. Wheeler. 2006. Phylogenetic systematics of Dart-poison Frogs and their relatives (Amphibia: Athesphatanura: Dendrobatidae). *Bull. Amer. Mus. Nat. Hist.* (299): 1–262.
- Gusovsky, F. and J.W. Daly. 1988. Formation of inositol phosphates in synaptoneurosomes of guinea pig brain: stimulatory effects of receptor agonists, sodium channel agents and sodium and calcium ionophores. *Neuropharmacology* 27:95–105.
- , E.B. Hollingsworth, and J.W. Daly. 1986. Regulation of phosphatidylinositol turnover in brain synaptoneurosomes: stimulatory effects of agents that enhance influx of sodium ions. *Proc. Natl. Acad. Sci. USA* 83:3003–3007.
- , E.T. McNeal, and J.W. Daly. 1987. Stimulation of phosphoinositide breakdown in brain synaptoneurosomes by agents that activate sodium influx: antagonism by tetrodotoxin, saxitoxin, and cadmium. *Mol. Pharmacol.* 32:479–487.
- , W.L. Padgett, C.R. Creveling, and J.W. Daly. 1992. Interaction of pumiliotoxin B with an “alkaloid-binding domain” on the voltage-dependent sodium channel. *Mol. Pharmacol.* 42:1104–1108.
- , D.P. Rossignol, E.T. McNeal, and J.W. Daly. 1988. Pumiliotoxin B binds to a site on the voltage-dependent sodium channel that is allosterically coupled to other binding sites. *Proc. Natl. Acad. Sci.* 85:1272–1276.
- Hagman, M. and A. Forsman. 2003. Correlated evolution of conspicuous coloration and body size in poison frogs (Dendrobatidae). *Evolution* 57: 2904–2910.
- Han, X. 2008. Does Life History Shape Sexual Size Dimorphism in Anurans: A Comparative Analysis. Ph.D. Diss., Univ. Guelph, Guelph, Ontario, Canada.
- Hödl, W. and A. Amezcuita. 2001. Visual signaling in anuran amphibians, p. 121–141. *In* M.J. Ryan (ed.), *Anuran Communication*. Smithsonian Inst. Press, Washington, D.C.
- Hollingsworth, E.B., E.B. Sears, R.A. de la Cruz, F. Gusovsky, and J.W. Daly. 1986. Accumulations of cyclic AMP and inositol phosphates in guinea pig cerebral cortical synaptoneurosomes: enhancement by agents acting at sodium channels. *Biochim. Biophys. Acta* 883:15–25.
- Jones, T.H., J.S.T. Gorman, R.R. Snelling, J.H.C. Delabie, M.S. Blum, H.M. Garraffo, P. Jain, J.W. Daly, and T.F. Spande. 1999. Further alkaloids common to ants and frogs: decahydroquinolines and a quinolizidine. *J. Chem. Ecol.* 25:1179–1193.
- Killeen, T.J. 2007. A perfect storm in the Amazon wilderness: development and conservation in the context of the initiative for the integration of the regional infrastructure of South America (IIRSA). *Conserv. Intl., Advances Appl. Biodivers. Sci.* (7): 1–99.
- Lahanas, P.N. 1992. Historical Biogeographic Relationships of Central and South America: A Biochemical, Phylogenetic Analysis of Selected Amphibians and Reptiles. Ph.D. Diss., Univ. Miami, Coral Gables, Florida.
- Laschat, S., F. Narjes, and L.E. Overman. 1994. Application of intramolecular Heck reactions to the preparation of steroid and terpene intermediates having cis A-B ring fusions. Model studies for the total synthesis of complex cardenolides. *Tetrahedron* 50:347–358.
- Lötters, S., F.C. Herrera, J. Kohler, and R. Richter. 1997. Notes on the distribution and color variation of poison frogs of the genus *Phyllobates* from western Colombia (Anura: Dendrobatidae). *Rev. Francaise d’aquariol. herpétol.* 24:55–58.
- Lovenberg, T. and J.W. Daly. 1986. Histronicotoxins: effects on binding of radioligands for sodium, potassium, and calcium channels in brain membranes. *Neurochem. Res.* 11:1609–1621.
- Martel, A., P. Van Rooij, G. Vercauteren, K. Baert, L. Van Waeyenberghe, P. Debacker, T.W.J. Garner, T. Woeltjes, R. Ducatelle, F. Haesebrouck, and F. Pasmans. 2011. Developing a safe antifungal treatment protocol to eliminate *Batrachochytrium dendrobatidis* from amphibians. *Med. Mycol.* 49: 143–149.
- Maxson, L.R. and C.W. Myers. 1985. Albumin evolution in tropical poison frogs (Dendrobatidae): a preliminary report. *Biotropica* 17:50–56.
- McCain, C.M. and N.J. Sanders. 2010. Metabolic theory and elevational diversity of vertebrate ectotherms. *Ecology* 91:601–609.
- Miller, D.L., S. Rajeev, M. Brookins, J. Cook, L. Whittington, and C.A. Baldwin. 2008. Concurrent infection with ranavirus, *Batrachochytrium dendrobatidis*, and *Aeromonas* in a captive anuran colony. *J. Zoo Wildl. Med.* 39:445–449.
- Moffett, M.W. 1995. Lurid and lethal: poison dart frogs. *Natl. Geogr. Mag.* 187:98–111.
- Müller, C.E. 1996. Epibatidin – ein nicotinartiges, analgetisch wirksames alkaloid aus pfeilgiftfröschen. *Pharm. Unserer Zeit* 25:85–92.
- Myers, C.W. 1982. Spotted Poison Frogs: descriptions of three new *Dendrobates* from western Amazonia, and resurrection of a lost species from “Chiriqui”. *Amer. Mus. Novitates* (2721):1–23.
- and J.W. Daly. 1983. Dart-poison frogs. *Sci. Amer.* 248:120–133.
- , –, and B. Malkin. 1978. A dangerously toxic new frog (*Phyllobates*) used by Emberá Indians of western Colombia, with discussion of blowgun fabrication and dart poisoning. *Bull. Amer. Mus. Nat. Hist.* 161:308–365.
- Neuwirth, M., J.W. Daly, C.W. Myers, and L.W. Tice. 1979. Morphology of the granular secretory glands in skin of poison-dart frogs (Dendrobatidae). *Tissue Cell* 11:755–771.
- Nijman, V. and C.R. Shepherd. 2010. The role of Asia in the global trade in CITES II-listed poison arrow frogs: hopping from Kazakhstan to Lebanon to Thailand and beyond. *Biodivers. Conserv.* 19: 1963–1970.
- Nogge, G. 2004. Jahresbericht 2003 der Aktiengesellschaft Zoologischer Garten Köln. *Z. Kölner*

- Zoo 47:1–32.
- Obst, F.J., K. Richter, and U. Jacob. 1988. The Completely Illustrated Atlas of Reptiles and Amphibians for the Terrarium. T.F.H Publ., Inc., Neptune City, New Jersey.
- Riston, J.R. 2006. Estudos Visando à Síntese Esterosseletiva do Alcalóide 275A. Diss. Mestrado, Univ. Estadual Campinas, Campinas, Brazil.
- Rueda-Almonacid, J.V. 1999. Anfíbios y reptiles amenazados de extinción en Colombia. Rev. Acad. Colombiana Cienc. Exact. Fís. Nat. 23:475–497.
- Ruiz-Carranza, P.M., M.C. Ardila-Robayo, and J.D. Lynch. 1996. Lista actualizada de la fauna de anfibios de Colombia. Rev. Acad. Colombiana Cienc. Exact. Fís. Nat. 20:365–415.
- Santos, J.C. 2009. Phylogeography and the Evolution of Correlated Traits under Multiple Origins of Aposematism in the Poison Frog Family. Ph.D. Diss., The University of Texas at Austin.
- Santos, J.C., L.A. Coloma, and D.C. Cannatella. 2003. Multiple, recurring origins of aposematism and diet specialization in poison frogs. Proc. Natl. Acad. Sci. USA 100:12792–12797.
- Steven, A. and L.E. Overman. 2007. Total synthesis of complex cyclotryptamine alkaloids: stereocontrolled construction of quaternary carbon stereocenters. Angew. Chem. Intl. Ed. 46:5488–5508.
- Tokuyama, T. and J.W. Daly. 1983. Steroidal alkaloids (batrachotoxins and 4 β -hydroxybatrachotoxins), “indole alkaloids” (calycanthine and chimonanthine) and a piperidinyldipyridin. Tetrahedron 39: 41–47.
- Toledo, L.F., R.S. Ribeiro, and C.F.B. Haddad. 2007. Anurans as prey: an exploratory analysis and size relationships between predators and their prey. J. Zool. 271:170–177.
- Widmer, A., S. Lötters, and K.-H. Jungfer. 2000. A molecular phylogenetic analysis of the Neotropical dart-poison frog genus *Phyllobates* (Amphibia: Dendrobatidae). Naturwissenschaften 87:559–562.
- Zimmermann, E. and H. Zimmermann. 1992. Dart-poison frogs (Dendrobatidae): biology; breeding and conservation, p. 5–20. In M.J. Uricheck (ed.), 15th International Herpetological Symposium on Captive Propagation & Husbandry. Intl. Herpetol. Symp., Inc.
- and –. 1994. Reproductive strategies, breeding, and conservation of tropical frogs: dart-poison frogs and Malagasy poison frogs, p. 255–266. In J.B. Murphy, K. Adler, and J.T. Collins (eds.), Captive Management and Conservation of Amphibians and Reptiles. SSAR Contrib. Herpetol. (11).
- Zimmermann, H. and E. Zimmermann. 1985. Zur Fortpflanzungsstrategie des Pfeilgiftfrosches *Phyllobates terribilis* Myers, Daly & Malkin, 1978. Salamandra 21:281–297.

Anthony M. Nowacki and **Tiffany M. Doan**, Central Connecticut State University, Department of Biology, 1615 Stanley Street, New Britain, CT 06050, USA (tony_nowacki@yahoo.com) and (tiffperu@yahoo.com).

Primary editor for this account, Andrew H. Price.

Published 30 April 2012 and Copyright © 2012 by the Society for the Study of Amphibians and Reptiles.
