

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

{ μ -6,6'-Dimethoxy-2,2'-[butane-1,4-diylbis(nitrilomethylidene)]diphenolato-1:2 κ^8 O⁶,O¹,O^{1'},O^{6'}:O¹,N,N',O^{1'}}tris-(nitrate-1 κ^2 O,O')copper(II)-gadolinium(III)

Christopher Chan, Xiaoping Yang, Richard A. Jones,*
Bradley J. Holliday* and Julie M. Stanley

Department of Chemistry & Biochemistry, The University of Texas at Austin,
1 University Station A5300, Austin, TX 78712-0165, USA
Correspondence e-mail: rajones@mail.utexas.edu, bholliday@cm.utexas.edu


Received 16 February 2010; accepted 21 April 2010

Key indicators: single-crystal X-ray study; $T = 223$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å;
 R factor = 0.033; wR factor = 0.128; data-to-parameter ratio = 12.3.

In the title dinuclear complex, $[\text{CuGd}(\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_4)(\text{NO}_3)_3]$, the Cu^{II} ion is located in the inner N_2O_2 cavity of the Schiff base ligand and adopts a distorted square-planar geometry. The Gd^{III} ion is ten-coordinate being bound to ten O atoms, four from the Schiff base ligand and six from three bidentate nitrate anions. The Cu^{II} and Gd^{III} ions are linked by two phenolate O atoms of the Schiff base ligand, with a separation of 3.5185 (9) Å.

Related literature

For general background to 3d-4f bimetallic complexes, see: Sakamoto *et al.* (2001); Winpenny (1998); Yang *et al.* (2005). For related structures, see: Fei & Fang (2008); Xing *et al.* (2008).



Experimental

Crystal data

$[\text{CuGd}(\text{C}_{20}\text{H}_{22}\text{N}_2\text{O}_4)(\text{NO}_3)_3]$
 $M_r = 761.22$
Monoclinic, $P2_1/n$
 $a = 11.795$ (2) Å
 $b = 14.730$ (3) Å
 $c = 14.892$ (3) Å
 $\beta = 100.58$ (3)°

$V = 2543.4$ (9) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 3.50$ mm⁻¹
 $T = 223$ K
 $0.56 \times 0.13 \times 0.13$ mm

Data collection

Rigaku MiniFlexII CCD
diffractometer
Absorption correction: multi-scan
(*CrystalClear*; Rigaku/MS,
2002)
 $T_{\text{min}} = 0.598$, $T_{\text{max}} = 1.000$

14367 measured reflections
4460 independent reflections
4090 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.128$
 $S = 1.20$
4460 reflections
362 parameters

6 restraints
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 2.69$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.35$ e Å⁻³

Table 1

Selected bond lengths (Å).

Gd1—O1	2.581 (4)	Gd1—O9	2.428 (4)
Gd1—O2	2.336 (3)	Gd1—O11	2.431 (4)
Gd1—O3	2.418 (3)	Gd1—O12	2.520 (3)
Gd1—O4	2.568 (3)	Cu1—O2	1.941 (3)
Gd1—O5	2.472 (5)	Cu1—O3	1.940 (3)
Gd1—O6	2.458 (5)	Cu1—N1	2.004 (4)
Gd1—O8	2.472 (4)	Cu1—N2	1.960 (4)

Data collection: *CrystalClear* (Rigaku/MS, 2002); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The authors acknowledge the Freshman Research Initiative, funded in part by the HHMI Undergraduate Science Education Award (52005907) and the National Science Foundation (CHE-0629136). Single crystal X-ray data were collected using instrumentation purchased with funds provided by the National Science Foundation (CHE-0741973). BJH wishes to acknowledge the Welch Foundation (F-1631), the Petroleum Research Fund administered by the American Chemical Society (47022-G3), the National Science Foundation (CHE-0639239, CHE-0847763), the American Heart Association (0765078Y), and the UT-CNM and UT-Austin for financial support of this research. RAJ wishes to acknowledge the Welch Foundation (F-816), the Texas Higher Education Coordinating Board (ARP003658-0010-2006) and the Petroleum Research Fund, administered by the American Chemical Society (47014-AC5).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2285).

References

- Fei, L. & Fang, Z. (2008). *Acta Cryst.* **E64**, m406.
- Rigaku/MS (2002). *CrystalClear*. Rigaku/MS, The Woodlands, Texas, USA.
- Sakamoto, M., Manseki, K. & Okawa, H. (2001). *Coord. Chem. Rev.* **219–221**, 379–414.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Winpenny, R. E. P. (1998). *Chem. Soc. Rev.* **27**, 447–452.
- Xing, J.-C., Wang, J.-H., Yan, P.-F. & Li, G.-M. (2008). *Acta Cryst.* **E64**, m1206.
- Yang, X., Jones, R. A., Lynch, V., Oye, M. M. & Holmes, A. L. (2005). *Dalton Trans.* pp. 849–851.

supplementary materials

Acta Cryst. (2010). E66, m576-m577 [doi:10.1107/S1600536810014716]

{ μ -6,6'-Dimethoxy-2,2'-[butane-1,4-diylbis(nitrilomethylidene)]diphenolato-1:2 κ^8 O⁶,O¹,O^{1'},O^{6'}:O¹,N,N',O^{1'}}tris(nitrato-1 κ^2 O,O')copper(II)gadolinium(III)

C. Chan, X. Yang, R. A. Jones, B. J. Holliday and J. M. Stanley

Comment

Heteropolynuclear complexes containing d- and f-block elements are currently of great interest because of their interesting physicochemical properties and potential applications as new materials (Sakamoto *et al.*, 2001; Winpenny, 1998). Compartmental Schiff bases with two dissimilar metal-binding sites, one being specific for the d-block metal ion and another for the f-block metal ion, are classical ligands used to synthesize such heteronuclear complexes. As part of our ongoing interests in 3d–4f complexes with Schiff-base ligands (Yang *et al.*, 2005), the title complex was synthesized and its crystal structure is reported herein.

The molecular structure of the title complex is shown in Fig. 1. It has a similar structure to other two Cu–Ln complexes with the Schiff-base ligand *N,N*-bis(3-methoxysalicylidene)propane-1,3-diamine (H₂L), [CuLnL(NO₃)₃] [Ln = Eu^{III} (Xing *et al.*, 2008) and Tb^{III} (Fei & Fang, 2008)]. The Cu^{II} ion is coordinated by two N atoms and two O atoms from the Schiff-base ligand. The Gd^{III} ion is surrounded by ten O atoms, four from the Schiff-base ligand and six from three bidentate NO₃[−] anions. Cu^{II} and Gd^{III} ions are bridged by two phenolate O atoms of the Schiff-base ligand, with a separation of 3.5185 (9) Å. The average distances for the Cu—O(phenolate) and Cu—N are 1.942 Å and 1.982 Å, respectively. The Gd—O(phenolate) distance of 2.377 Å (*av.*) is shorter than the Gd—O (methoxy) distance of 2.574 Å (*av.*), no doubt reflecting the difference between ionic vs. dative bonding. The average distance for the Gd—O (nitrate) is 2.464 Å, which is comparable to those found in the literature (Fei & Fang, 2008; Xing *et al.*, 2008).

Experimental

A mixture of the Schiff-base ligand (0.178 g, 0.5 mmol) and Cu(CH₃CO₂)₂·H₂O (0.10 g, 0.5 mmol) in EtOH (15 ml) was stirred and refluxed for 10 min. The reaction mixture was allowed to cool briefly and Gd(NO₃)₃·6H₂O (0.226 g, 0.5 mmol) was added and the mixture again heated under reflux (15 min) and then filtered. Diethylether was allowed to diffuse slowly into the filtrate at room temperature and blue crystals were obtained after two weeks. The crystals were filtered off and washed with 5 ml of EtOH (yield 0.176 g, 46.32%). m. p. > 260°C (dec). ESI-MS(MeOH) *m/z*: 700 [M—NO₃]⁺. IR(CH₃OH, cm^{−1}): 3429 m, 1636 s, 1473 s, 1439 w, 1362 m, 1295 w, 1227 m, 1171 w, 1099 w, 1077 m, 1013 m, 658 s. UV-VIS(CH₃OH, 25°C) λ_{\max} /nm: 215, 275, 350.

Refinement

All H atoms were positioned geometrically and refined using a riding model, with C—H = 0.94 (CH), 0.98 (CH₂) and 0.97 (CH₃) Å and with $U_{\text{iso}}(\text{H}) = 1.2(1.5 \text{ for methyl groups})U_{\text{eq}}(\text{C})$. The highest residual electron density was found 1.83 Å from H1A and the deepest hole 0.72 Å from Cu1.

Figures




Fig. 1. The molecular structure of the title complex, with displacement ellipsoids shown at the 40% probability level. H atoms have been excluded for clarity.

{ μ -6,6'-Dimethoxy-2,2'-[butane-1,4-diylbis(nitrilomethylidene)]diphenolato- 1:2 $\kappa^8 O^6, O^1, O^1, O^6$:
 O^1, N, N^1, O^1 }tris(nitrato- 1 $\kappa^2 O, O^1$)copper(II)gadolinium(III)

Crystal data

[CuGd(C₂₀H₂₂N₂O₄)(NO₃)₃]

$M_r = 761.22$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 11.795$ (2) Å

$b = 14.730$ (3) Å

$c = 14.892$ (3) Å

$\beta = 100.58$ (3)°

$V = 2543.4$ (9) Å³

$Z = 4$

$F(000) = 1496$

$D_x = 1.988$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 8776 reflections

$\theta = 1.8$ – 31.9 °

$\mu = 3.50$ mm⁻¹

$T = 223$ K

Prism, blue

$0.56 \times 0.13 \times 0.13$ mm

Data collection

Rigaku MiniFlexII CCD
 diffractometer

Radiation source: fine-focus sealed tube
 graphite

Detector resolution: 13.6612 pixels mm⁻¹

thin-slice ω scans

Absorption correction: multi-scan
 (CrystalClear; Rigaku/MS, 2002)

$T_{\min} = 0.598$, $T_{\max} = 1.000$

14367 measured reflections

4460 independent reflections

4090 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.032$

$\theta_{\max} = 25.0$ °, $\theta_{\min} = 2.0$ °

$h = -14 \rightarrow 14$

$k = -17 \rightarrow 17$

$l = -17 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.033$

$wR(F^2) = 0.128$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring
 sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0793P)^2 + 1.3271P]$

$S = 1.20$

4460 reflections

362 parameters

6 restraints

Primary atom site location: structure-invariant direct methods

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 2.69 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -1.35 \text{ e } \text{\AA}^{-3}$

Extinction correction: SHELXTL (Sheldrick, 2008)

Extinction coefficient: 0.0022 (4)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Gd1	0.261236 (19)	0.750997 (12)	0.066103 (16)	0.02180 (16)
Cu1	0.54871 (5)	0.76924 (4)	0.03441 (4)	0.02328 (19)
N1	0.6472 (3)	0.6645 (3)	0.0102 (3)	0.0281 (9)
N2	0.6638 (3)	0.8666 (3)	0.0545 (3)	0.0251 (8)
O1	0.2007 (3)	0.6647 (3)	-0.0860 (2)	0.0345 (8)
O2	0.4094 (3)	0.6988 (2)	-0.0053 (2)	0.0305 (7)
O3	0.4328 (3)	0.8431 (2)	0.0767 (2)	0.0275 (7)
O4	0.2582 (3)	0.9074 (2)	0.1417 (2)	0.0337 (8)
C1	0.0855 (5)	0.6569 (5)	-0.1376 (5)	0.0605 (18)
H1A	0.0850	0.6743	-0.2005	0.091*
H1B	0.0345	0.6967	-0.1116	0.091*
H1C	0.0593	0.5947	-0.1356	0.091*
C2	0.2860 (4)	0.6140 (3)	-0.1153 (3)	0.0294 (10)
C3	0.2659 (5)	0.5475 (3)	-0.1817 (3)	0.0345 (11)
H3A	0.1904	0.5355	-0.2124	0.041*
C4	0.3593 (5)	0.4980 (4)	-0.2032 (4)	0.0406 (13)
H4A	0.3463	0.4524	-0.2482	0.049*
C5	0.4697 (5)	0.5162 (4)	-0.1584 (4)	0.0386 (12)
H5A	0.5316	0.4820	-0.1721	0.046*
C6	0.4913 (4)	0.5862 (3)	-0.0915 (3)	0.0306 (10)
C7	0.3977 (4)	0.6343 (3)	-0.0690 (3)	0.0277 (10)
C8	0.6085 (4)	0.5997 (3)	-0.0437 (3)	0.0326 (11)
H8A	0.6625	0.5556	-0.0536	0.039*
C9	0.7715 (4)	0.6524 (4)	0.0540 (4)	0.0402 (14)
H9A	0.8203	0.6674	0.0095	0.048*
H9B	0.7850	0.5886	0.0713	0.048*
C10	0.8059 (5)	0.7113 (5)	0.1378 (4)	0.0426 (13)
H10A	0.8640	0.6792	0.1820	0.051*
H10B	0.7383	0.7215	0.1662	0.051*
C11	0.8552 (4)	0.8039 (4)	0.1151 (4)	0.0433 (14)
H11A	0.8623	0.8434	0.1688	0.052*
H11B	0.9327	0.7948	0.1016	0.052*
C12	0.7812 (5)	0.8503 (4)	0.0350 (4)	0.0338 (12)
H12A	0.8164	0.9083	0.0230	0.041*
H12B	0.7760	0.8123	-0.0196	0.041*
C13	0.6435 (4)	0.9476 (3)	0.0807 (3)	0.0263 (10)
H13A	0.7029	0.9901	0.0806	0.032*

supplementary materials

C14	0.5420 (4)	0.9812 (3)	0.1102 (3)	0.0238 (9)
C15	0.5441 (4)	1.0721 (3)	0.1420 (3)	0.0286 (10)
H15A	0.6097	1.1082	0.1417	0.034*
C16	0.4500 (4)	1.1079 (4)	0.1733 (3)	0.0349 (11)
H16A	0.4512	1.1684	0.1937	0.042*
C17	0.3544 (4)	1.0541 (3)	0.1745 (3)	0.0307 (10)
H17A	0.2913	1.0779	0.1972	0.037*
C18	0.3500 (4)	0.9666 (3)	0.1433 (3)	0.0261 (9)
C19	0.4434 (4)	0.9281 (3)	0.1088 (3)	0.0219 (9)
C20	0.1675 (5)	0.9375 (5)	0.1878 (5)	0.0593 (18)
H20A	0.2008	0.9574	0.2491	0.089*
H20B	0.1146	0.8877	0.1912	0.089*
H20C	0.1262	0.9875	0.1542	0.089*
N3	0.3210 (8)	0.7319 (5)	0.2617 (4)	0.085 (3)
O5	0.3967 (6)	0.7260 (5)	0.2116 (4)	0.0811 (19)
O6	0.2209 (7)	0.7391 (3)	0.2219 (4)	0.0643 (17)
O7	0.3486 (8)	0.7274 (6)	0.3444 (5)	0.135 (3)
N4	0.1158 (4)	0.8781 (3)	-0.0484 (3)	0.0408 (11)
O8	0.2183 (3)	0.8626 (3)	-0.0595 (2)	0.0370 (9)
O9	0.0797 (3)	0.8273 (3)	0.0094 (3)	0.0380 (8)
O10	0.0586 (5)	0.9393 (4)	-0.0899 (4)	0.0827 (17)
N5	0.1497 (4)	0.5795 (3)	0.0913 (3)	0.0397 (10)
O11	0.2586 (3)	0.5895 (3)	0.0994 (3)	0.0396 (9)
O12	0.0899 (3)	0.6504 (3)	0.0719 (3)	0.0421 (9)
O13	0.1069 (5)	0.5050 (3)	0.1020 (3)	0.0727 (15)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Gd1	0.0171 (2)	0.0226 (2)	0.0268 (2)	-0.00134 (7)	0.00685 (14)	-0.00039 (7)
Cu1	0.0182 (3)	0.0237 (3)	0.0291 (3)	-0.0009 (2)	0.0074 (2)	-0.0030 (2)
N1	0.0213 (19)	0.023 (2)	0.042 (2)	0.0040 (16)	0.0126 (17)	0.0029 (18)
N2	0.0184 (18)	0.028 (2)	0.031 (2)	-0.0008 (16)	0.0101 (15)	-0.0004 (17)
O1	0.0271 (18)	0.037 (2)	0.0385 (18)	-0.0008 (16)	0.0028 (15)	-0.0079 (17)
O2	0.0229 (16)	0.0314 (18)	0.0381 (18)	-0.0031 (14)	0.0085 (14)	-0.0118 (15)
O3	0.0225 (16)	0.0221 (16)	0.0408 (18)	-0.0048 (13)	0.0132 (14)	-0.0067 (14)
O4	0.0230 (16)	0.0337 (19)	0.049 (2)	-0.0025 (15)	0.0196 (15)	-0.0092 (16)
C1	0.029 (3)	0.078 (5)	0.067 (4)	-0.001 (3)	-0.010 (3)	-0.026 (4)
C2	0.027 (2)	0.030 (3)	0.031 (2)	0.000 (2)	0.0073 (19)	0.001 (2)
C3	0.040 (3)	0.033 (3)	0.028 (2)	-0.008 (2)	0.002 (2)	-0.005 (2)
C4	0.053 (3)	0.037 (3)	0.035 (3)	-0.001 (3)	0.016 (2)	-0.009 (2)
C5	0.049 (3)	0.029 (3)	0.043 (3)	-0.002 (2)	0.022 (2)	-0.005 (2)
C6	0.035 (3)	0.024 (2)	0.035 (3)	-0.003 (2)	0.014 (2)	-0.002 (2)
C7	0.035 (3)	0.022 (2)	0.028 (2)	-0.0046 (19)	0.0119 (19)	-0.0014 (19)
C8	0.031 (2)	0.025 (2)	0.046 (3)	0.004 (2)	0.017 (2)	-0.003 (2)
C9	0.022 (3)	0.033 (3)	0.066 (4)	0.006 (2)	0.007 (3)	0.007 (3)
C10	0.026 (3)	0.058 (4)	0.043 (3)	0.004 (3)	0.004 (2)	0.010 (3)
C11	0.019 (2)	0.057 (4)	0.054 (3)	0.000 (2)	0.007 (2)	-0.013 (3)

C12	0.028 (3)	0.030 (3)	0.050 (3)	-0.005 (2)	0.024 (2)	-0.002 (2)
C13	0.026 (2)	0.027 (2)	0.027 (2)	-0.0076 (19)	0.0066 (18)	0.0012 (19)
C14	0.025 (2)	0.024 (2)	0.023 (2)	0.0006 (18)	0.0039 (17)	-0.0005 (18)
C15	0.031 (2)	0.025 (2)	0.030 (2)	-0.002 (2)	0.0064 (19)	-0.005 (2)
C16	0.034 (3)	0.029 (3)	0.041 (3)	0.002 (2)	0.004 (2)	-0.008 (2)
C17	0.026 (2)	0.031 (2)	0.038 (2)	0.004 (2)	0.0113 (19)	-0.006 (2)
C18	0.026 (2)	0.026 (2)	0.028 (2)	-0.0019 (18)	0.0096 (18)	-0.0008 (19)
C19	0.021 (2)	0.022 (2)	0.023 (2)	0.0007 (18)	0.0052 (16)	-0.0005 (18)
C20	0.041 (3)	0.065 (4)	0.083 (5)	-0.008 (3)	0.041 (3)	-0.030 (4)
N3	0.108 (7)	0.111 (5)	0.032 (3)	-0.088 (5)	-0.002 (4)	0.011 (3)
O5	0.068 (4)	0.113 (4)	0.052 (3)	-0.045 (4)	-0.015 (3)	0.027 (3)
O6	0.099 (5)	0.056 (3)	0.047 (3)	-0.031 (3)	0.037 (3)	-0.004 (2)
O7	0.149 (6)	0.203 (7)	0.047 (3)	-0.114 (5)	0.000 (4)	0.020 (4)
N4	0.038 (2)	0.036 (3)	0.048 (3)	0.001 (2)	0.008 (2)	0.006 (2)
O8	0.0287 (18)	0.035 (2)	0.049 (2)	-0.0025 (16)	0.0136 (16)	0.0100 (17)
O9	0.0300 (18)	0.041 (2)	0.044 (2)	-0.0057 (16)	0.0112 (16)	0.0052 (17)
O10	0.081 (4)	0.066 (3)	0.102 (4)	0.030 (3)	0.019 (3)	0.048 (3)
N5	0.044 (3)	0.034 (2)	0.040 (2)	-0.016 (2)	0.006 (2)	-0.001 (2)
O11	0.037 (2)	0.0297 (19)	0.053 (2)	0.0049 (16)	0.0118 (17)	0.0050 (18)
O12	0.0273 (19)	0.042 (2)	0.057 (2)	-0.0071 (17)	0.0080 (17)	0.0070 (19)
O13	0.099 (4)	0.041 (3)	0.078 (3)	-0.028 (3)	0.014 (3)	0.007 (2)

Geometric parameters (Å, °)

Gd1—O1	2.581 (4)	C6—C8	1.447 (7)
Gd1—O2	2.336 (3)	C8—H8A	0.9400
Gd1—O3	2.418 (3)	C9—C10	1.512 (9)
Gd1—O4	2.568 (3)	C9—H9A	0.9800
Gd1—O5	2.472 (5)	C9—H9B	0.9800
Gd1—O6	2.458 (5)	C10—C11	1.544 (9)
Gd1—O8	2.472 (4)	C10—H10A	0.9800
Gd1—O9	2.428 (4)	C10—H10B	0.9800
Gd1—O11	2.431 (4)	C11—C12	1.507 (8)
Gd1—O12	2.520 (3)	C11—H11A	0.9800
Cu1—O2	1.941 (3)	C11—H11B	0.9800
Cu1—O3	1.940 (3)	C12—H12A	0.9800
Cu1—N1	2.004 (4)	C12—H12B	0.9800
Cu1—N2	1.960 (4)	C13—C14	1.436 (6)
N1—C8	1.276 (6)	C13—H13A	0.9400
N1—C9	1.502 (6)	C14—C19	1.398 (6)
N2—C13	1.292 (6)	C14—C15	1.419 (6)
N2—C12	1.486 (6)	C15—C16	1.385 (7)
O1—C2	1.386 (6)	C15—H15A	0.9400
O1—C1	1.437 (6)	C16—C17	1.382 (7)
O2—C7	1.332 (6)	C16—H16A	0.9400
O3—C19	1.338 (5)	C17—C18	1.367 (7)
O4—C18	1.387 (6)	C17—H17A	0.9400
O4—C20	1.443 (6)	C18—C19	1.417 (6)
C1—H1A	0.9700	C20—H20A	0.9700

supplementary materials

C1—H1B	0.9700	C20—H20B	0.9700
C1—H1C	0.9700	C20—H20C	0.9700
C2—C3	1.381 (7)	N3—O7	1.218 (9)
C2—C7	1.403 (7)	N3—O6	1.224 (11)
C3—C4	1.406 (7)	N3—O5	1.266 (11)
C3—H3A	0.9400	N4—O10	1.224 (6)
C4—C5	1.377 (8)	N4—O8	1.270 (6)
C4—H4A	0.9400	N4—O9	1.272 (5)
C5—C6	1.424 (7)	N5—O13	1.230 (6)
C5—H5A	0.9400	N5—O12	1.264 (6)
C6—C7	1.404 (7)	N5—O11	1.277 (6)
O2—Gd1—O3	61.77 (11)	C2—C3—H3A	120.2
O2—Gd1—O9	132.81 (12)	C4—C3—H3A	120.2
O3—Gd1—O9	115.90 (12)	C5—C4—C3	119.9 (5)
O2—Gd1—O11	79.02 (12)	C5—C4—H4A	120.0
O3—Gd1—O11	125.29 (12)	C3—C4—H4A	120.0
O9—Gd1—O11	118.72 (12)	C4—C5—C6	120.8 (5)
O2—Gd1—O6	134.1 (2)	C4—C5—H5A	119.6
O3—Gd1—O6	106.44 (17)	C6—C5—H5A	119.6
O9—Gd1—O6	92.9 (2)	C7—C6—C5	119.0 (5)
O11—Gd1—O6	74.10 (14)	C7—C6—C8	122.4 (4)
O2—Gd1—O8	86.60 (12)	C5—C6—C8	118.4 (5)
O3—Gd1—O8	74.26 (12)	O2—C7—C2	117.9 (4)
O9—Gd1—O8	51.97 (12)	O2—C7—C6	123.2 (4)
O11—Gd1—O8	142.79 (13)	C2—C7—C6	119.0 (4)
O6—Gd1—O8	135.51 (17)	N1—C8—C6	127.7 (4)
O2—Gd1—O5	86.1 (2)	N1—C8—H8A	116.2
O3—Gd1—O5	68.18 (15)	C6—C8—H8A	116.2
O9—Gd1—O5	139.5 (2)	N1—C9—C10	112.6 (5)
O11—Gd1—O5	72.88 (18)	N1—C9—H9A	109.1
O6—Gd1—O5	50.8 (2)	C10—C9—H9A	109.1
O8—Gd1—O5	140.55 (16)	N1—C9—H9B	109.1
O2—Gd1—O12	119.60 (12)	C10—C9—H9B	109.1
O3—Gd1—O12	174.12 (12)	H9A—C9—H9B	107.8
O9—Gd1—O12	67.83 (12)	C9—C10—C11	112.4 (5)
O11—Gd1—O12	51.51 (13)	C9—C10—H10A	109.1
O6—Gd1—O12	68.33 (17)	C11—C10—H10A	109.1
O8—Gd1—O12	111.26 (12)	C9—C10—H10B	109.1
O5—Gd1—O12	105.99 (16)	C11—C10—H10B	109.1
O2—Gd1—O4	124.51 (11)	H10A—C10—H10B	107.9
O3—Gd1—O4	63.05 (10)	C12—C11—C10	112.8 (4)
O9—Gd1—O4	69.79 (12)	C12—C11—H11A	109.0
O11—Gd1—O4	141.89 (12)	C10—C11—H11A	109.0
O6—Gd1—O4	68.24 (13)	C12—C11—H11B	109.0
O8—Gd1—O4	73.61 (12)	C10—C11—H11B	109.0
O5—Gd1—O4	78.95 (19)	H11A—C11—H11B	107.8
O12—Gd1—O4	115.88 (12)	N2—C12—C11	110.1 (4)
O2—Gd1—O1	63.16 (11)	N2—C12—H12A	109.6
O3—Gd1—O1	115.29 (11)	C11—C12—H12A	109.6

O9—Gd1—O1	80.71 (12)	N2—C12—H12B	109.6
O11—Gd1—O1	71.65 (13)	C11—C12—H12B	109.6
O6—Gd1—O1	136.28 (15)	H12A—C12—H12B	108.2
O8—Gd1—O1	71.23 (12)	N2—C13—C14	128.4 (4)
O5—Gd1—O1	136.5 (2)	N2—C13—H13A	115.8
O12—Gd1—O1	69.26 (13)	C14—C13—H13A	115.8
O4—Gd1—O1	143.41 (12)	C19—C14—C15	119.7 (4)
O3—Cu1—O2	77.94 (13)	C19—C14—C13	122.7 (4)
O3—Cu1—N2	92.82 (14)	C15—C14—C13	117.6 (4)
O2—Cu1—N2	164.02 (16)	C16—C15—C14	120.5 (5)
O3—Cu1—N1	163.09 (15)	C16—C15—H15A	119.8
O2—Cu1—N1	91.16 (15)	C14—C15—H15A	119.8
N2—Cu1—N1	100.54 (16)	C17—C16—C15	119.4 (5)
O3—Cu1—Gd1	41.03 (9)	C17—C16—H16A	120.3
O2—Cu1—Gd1	38.50 (9)	C15—C16—H16A	120.3
N2—Cu1—Gd1	133.86 (11)	C18—C17—C16	121.1 (4)
N1—Cu1—Gd1	124.78 (11)	C18—C17—H17A	119.4
C8—N1—C9	113.1 (4)	C16—C17—H17A	119.4
C8—N1—Cu1	122.5 (3)	C17—C18—O4	125.4 (4)
C9—N1—Cu1	124.4 (3)	C17—C18—C19	121.2 (4)
C13—N2—C12	116.1 (4)	O4—C18—C19	113.5 (4)
C13—N2—Cu1	124.1 (3)	O3—C19—C14	123.5 (4)
C12—N2—Cu1	119.7 (3)	O3—C19—C18	118.5 (4)
C2—O1—C1	117.3 (4)	C14—C19—C18	118.1 (4)
C2—O1—Gd1	116.2 (3)	O4—C20—H20A	109.5
C1—O1—Gd1	126.2 (3)	O4—C20—H20B	109.5
C7—O2—Cu1	124.7 (3)	H20A—C20—H20B	109.5
C7—O2—Gd1	124.8 (3)	O4—C20—H20C	109.5
Cu1—O2—Gd1	110.36 (14)	H20A—C20—H20C	109.5
C19—O3—Cu1	128.0 (3)	H20B—C20—H20C	109.5
C19—O3—Gd1	124.8 (3)	O7—N3—O6	123.3 (10)
Cu1—O3—Gd1	107.17 (14)	O7—N3—O5	120.4 (10)
C18—O4—C20	116.3 (4)	O6—N3—O5	116.2 (6)
C18—O4—Gd1	119.9 (3)	N3—O5—Gd1	95.4 (5)
C20—O4—Gd1	123.7 (3)	N3—O6—Gd1	97.3 (5)
O1—C1—H1A	109.5	O10—N4—O8	121.1 (5)
O1—C1—H1B	109.5	O10—N4—O9	123.6 (5)
H1A—C1—H1B	109.5	O8—N4—O9	115.3 (4)
O1—C1—H1C	109.5	N4—O8—Gd1	95.1 (3)
H1A—C1—H1C	109.5	N4—O9—Gd1	97.2 (3)
H1B—C1—H1C	109.5	O13—N5—O12	122.9 (5)
C3—C2—O1	124.7 (4)	O13—N5—O11	121.3 (5)
C3—C2—C7	121.7 (4)	O12—N5—O11	115.8 (4)
O1—C2—C7	113.6 (4)	N5—O11—Gd1	98.3 (3)
C2—C3—C4	119.5 (5)	N5—O12—Gd1	94.4 (3)
O2—Gd1—Cu1—O3	159.0 (2)	O12—Gd1—O4—C18	178.8 (3)
O9—Gd1—Cu1—O3	59.78 (19)	O1—Gd1—O4—C18	-91.6 (4)
O11—Gd1—Cu1—O3	-149.68 (18)	N4—Gd1—O4—C18	-102.1 (3)
O6—Gd1—Cu1—O3	-73.1 (2)	N3—Gd1—O4—C18	101.7 (4)

supplementary materials

O8—Gd1—Cu1—O3	67.60 (18)	O2—Gd1—O4—C20	-178.1 (4)
O5—Gd1—Cu1—O3	-81.8 (2)	O3—Gd1—O4—C20	-171.7 (5)
O12—Gd1—Cu1—O3	-168.9 (2)	O9—Gd1—O4—C20	53.2 (5)
O4—Gd1—Cu1—O3	-5.48 (18)	O11—Gd1—O4—C20	-58.0 (5)
O1—Gd1—Cu1—O3	138.55 (18)	O6—Gd1—O4—C20	-48.6 (5)
N4—Gd1—Cu1—O3	62.47 (19)	O8—Gd1—O4—C20	108.0 (5)
N3—Gd1—Cu1—O3	-76.2 (2)	O5—Gd1—O4—C20	-100.7 (5)
O3—Gd1—Cu1—O2	-159.0 (2)	O12—Gd1—O4—C20	1.9 (5)
O9—Gd1—Cu1—O2	-99.2 (2)	O1—Gd1—O4—C20	91.5 (5)
O11—Gd1—Cu1—O2	51.3 (2)	N4—Gd1—O4—C20	81.0 (5)
O6—Gd1—Cu1—O2	127.9 (2)	N3—Gd1—O4—C20	-75.2 (5)
O8—Gd1—Cu1—O2	-91.38 (19)	C1—O1—C2—C3	-9.8 (8)
O5—Gd1—Cu1—O2	119.2 (2)	Gd1—O1—C2—C3	164.2 (4)
O12—Gd1—Cu1—O2	32.1 (2)	C1—O1—C2—C7	172.0 (5)
O4—Gd1—Cu1—O2	-164.46 (19)	Gd1—O1—C2—C7	-14.0 (5)
O1—Gd1—Cu1—O2	-20.43 (19)	O1—C2—C3—C4	-177.4 (5)
N4—Gd1—Cu1—O2	-96.5 (2)	C7—C2—C3—C4	0.7 (8)
N3—Gd1—Cu1—O2	124.9 (2)	C2—C3—C4—C5	-0.3 (8)
O2—Gd1—Cu1—N2	159.0 (2)	C3—C4—C5—C6	-1.3 (8)
O3—Gd1—Cu1—N2	0.0 (2)	C4—C5—C6—C7	2.5 (8)
O9—Gd1—Cu1—N2	59.8 (2)	C4—C5—C6—C8	177.4 (5)
O11—Gd1—Cu1—N2	-149.64 (19)	Cu1—O2—C7—C2	-155.9 (3)
O6—Gd1—Cu1—N2	-73.1 (2)	Gd1—O2—C7—C2	19.0 (6)
O8—Gd1—Cu1—N2	67.64 (18)	Cu1—O2—C7—C6	24.1 (6)
O5—Gd1—Cu1—N2	-81.7 (2)	Gd1—O2—C7—C6	-160.9 (3)
O12—Gd1—Cu1—N2	-168.8 (2)	C3—C2—C7—O2	-179.4 (4)
O4—Gd1—Cu1—N2	-5.44 (18)	O1—C2—C7—O2	-1.1 (6)
O1—Gd1—Cu1—N2	138.59 (18)	C3—C2—C7—C6	0.5 (7)
N4—Gd1—Cu1—N2	62.51 (19)	O1—C2—C7—C6	178.8 (4)
N3—Gd1—Cu1—N2	-76.1 (2)	C5—C6—C7—O2	177.8 (4)
O2—Gd1—Cu1—N1	-33.5 (2)	C8—C6—C7—O2	3.2 (7)
O3—Gd1—Cu1—N1	167.5 (2)	C5—C6—C7—C2	-2.1 (7)
O9—Gd1—Cu1—N1	-132.7 (2)	C8—C6—C7—C2	-176.8 (4)
O11—Gd1—Cu1—N1	17.83 (18)	C9—N1—C8—C6	174.9 (5)
O6—Gd1—Cu1—N1	94.39 (19)	Cu1—N1—C8—C6	-3.7 (7)
O8—Gd1—Cu1—N1	-124.89 (17)	C7—C6—C8—N1	-14.0 (8)
O5—Gd1—Cu1—N1	85.7 (2)	C5—C6—C8—N1	171.3 (5)
O12—Gd1—Cu1—N1	-1.4 (2)	C8—N1—C9—C10	-162.0 (5)
O4—Gd1—Cu1—N1	162.03 (17)	Cu1—N1—C9—C10	16.6 (7)
O1—Gd1—Cu1—N1	-53.95 (17)	N1—C9—C10—C11	-92.0 (6)
N4—Gd1—Cu1—N1	-130.02 (18)	C9—C10—C11—C12	48.1 (7)
N3—Gd1—Cu1—N1	91.3 (2)	C13—N2—C12—C11	99.6 (5)
O3—Cu1—N1—C8	70.4 (7)	Cu1—N2—C12—C11	-83.1 (5)
O2—Cu1—N1—C8	21.1 (4)	C10—C11—C12—N2	59.9 (6)
N2—Cu1—N1—C8	-147.9 (4)	C12—N2—C13—C14	-175.3 (5)
Gd1—Cu1—N1—C8	41.2 (4)	Cu1—N2—C13—C14	7.6 (7)
O3—Cu1—N1—C9	-108.1 (6)	N2—C13—C14—C19	-4.1 (7)
O2—Cu1—N1—C9	-157.4 (4)	N2—C13—C14—C15	175.6 (5)
N2—Cu1—N1—C9	33.6 (4)	C19—C14—C15—C16	1.3 (7)

Gd1—Cu1—N1—C9	-137.3 (4)	C13—C14—C15—C16	-178.5 (4)
O3—Cu1—N2—C13	-4.2 (4)	C14—C15—C16—C17	0.8 (7)
O2—Cu1—N2—C13	49.9 (7)	C15—C16—C17—C18	-1.5 (7)
N1—Cu1—N2—C13	-173.7 (4)	C16—C17—C18—O4	180.0 (5)
Gd1—Cu1—N2—C13	-4.2 (5)	C16—C17—C18—C19	0.2 (7)
O3—Cu1—N2—C12	178.8 (4)	C20—O4—C18—C17	-8.6 (7)
O2—Cu1—N2—C12	-127.2 (5)	Gd1—O4—C18—C17	174.3 (4)
N1—Cu1—N2—C12	9.2 (4)	C20—O4—C18—C19	171.2 (5)
Gd1—Cu1—N2—C12	178.8 (3)	Gd1—O4—C18—C19	-5.9 (5)
O2—Gd1—O1—C2	16.4 (3)	Cu1—O3—C19—C14	5.4 (6)
O3—Gd1—O1—C2	50.7 (3)	Gd1—O3—C19—C14	-176.9 (3)
O9—Gd1—O1—C2	165.1 (3)	Cu1—O3—C19—C18	-174.7 (3)
O11—Gd1—O1—C2	-70.4 (3)	Gd1—O3—C19—C18	2.9 (5)
O6—Gd1—O1—C2	-110.5 (4)	C15—C14—C19—O3	177.3 (4)
O8—Gd1—O1—C2	112.2 (3)	C13—C14—C19—O3	-3.0 (7)
O5—Gd1—O1—C2	-33.6 (4)	C15—C14—C19—C18	-2.5 (6)
O12—Gd1—O1—C2	-125.2 (3)	C13—C14—C19—C18	177.2 (4)
O4—Gd1—O1—C2	128.9 (3)	C17—C18—C19—O3	-178.0 (4)
N4—Gd1—O1—C2	138.9 (3)	O4—C18—C19—O3	2.2 (6)
N3—Gd1—O1—C2	-73.1 (5)	C17—C18—C19—C14	1.9 (7)
O2—Gd1—O1—C1	-170.3 (5)	O4—C18—C19—C14	-178.0 (4)
O3—Gd1—O1—C1	-135.9 (5)	O7—N3—O5—Gd1	-177.0 (8)
O9—Gd1—O1—C1	-21.6 (5)	O6—N3—O5—Gd1	5.4 (7)
O11—Gd1—O1—C1	103.0 (5)	O2—Gd1—O5—N3	-166.0 (5)
O6—Gd1—O1—C1	62.8 (6)	O3—Gd1—O5—N3	132.9 (5)
O8—Gd1—O1—C1	-74.5 (5)	O9—Gd1—O5—N3	28.1 (6)
O5—Gd1—O1—C1	139.7 (5)	O11—Gd1—O5—N3	-86.3 (5)
O12—Gd1—O1—C1	48.1 (5)	O6—Gd1—O5—N3	-3.1 (4)
O4—Gd1—O1—C1	-57.8 (5)	O8—Gd1—O5—N3	114.1 (4)
N4—Gd1—O1—C1	-47.8 (5)	O12—Gd1—O5—N3	-46.3 (5)
N3—Gd1—O1—C1	100.3 (6)	O4—Gd1—O5—N3	67.7 (5)
O3—Cu1—O2—C7	161.6 (4)	O1—Gd1—O5—N3	-122.8 (5)
N2—Cu1—O2—C7	105.9 (6)	N4—Gd1—O5—N3	70.5 (5)
N1—Cu1—O2—C7	-31.4 (4)	O7—N3—O6—Gd1	177.0 (8)
Gd1—Cu1—O2—C7	175.6 (5)	O5—N3—O6—Gd1	-5.5 (8)
O3—Cu1—O2—Gd1	-13.93 (15)	O2—Gd1—O6—N3	27.4 (5)
N2—Cu1—O2—Gd1	-69.7 (6)	O3—Gd1—O6—N3	-39.0 (5)
N1—Cu1—O2—Gd1	153.03 (17)	O9—Gd1—O6—N3	-157.0 (4)
O3—Gd1—O2—C7	-163.2 (4)	O11—Gd1—O6—N3	83.9 (5)
O9—Gd1—O2—C7	-62.9 (4)	O8—Gd1—O6—N3	-123.0 (5)
O11—Gd1—O2—C7	56.3 (4)	O5—Gd1—O6—N3	3.2 (5)
O6—Gd1—O2—C7	111.0 (4)	O12—Gd1—O6—N3	138.2 (5)
O8—Gd1—O2—C7	-89.2 (4)	O4—Gd1—O6—N3	-90.2 (5)
O5—Gd1—O2—C7	129.6 (4)	O1—Gd1—O6—N3	123.4 (4)
O12—Gd1—O2—C7	23.4 (4)	N4—Gd1—O6—N3	-141.5 (4)
O4—Gd1—O2—C7	-156.6 (3)	O10—N4—O8—Gd1	171.6 (5)
O1—Gd1—O2—C7	-18.6 (3)	O9—N4—O8—Gd1	-6.9 (5)
N4—Gd1—O2—C7	-79.8 (4)	O2—Gd1—O8—N4	159.8 (3)
N3—Gd1—O2—C7	123.0 (4)	O3—Gd1—O8—N4	-138.6 (3)

supplementary materials

O3—Gd1—O2—Cu1	12.39 (14)	O9—Gd1—O8—N4	4.2 (3)
O9—Gd1—O2—Cu1	112.66 (18)	O11—Gd1—O8—N4	92.9 (3)
O11—Gd1—O2—Cu1	-128.16 (19)	O6—Gd1—O8—N4	-41.0 (4)
O6—Gd1—O2—Cu1	-73.4 (2)	O5—Gd1—O8—N4	-120.5 (4)
O8—Gd1—O2—Cu1	86.33 (17)	O12—Gd1—O8—N4	39.2 (3)
O5—Gd1—O2—Cu1	-54.9 (2)	O4—Gd1—O8—N4	-72.7 (3)
O12—Gd1—O2—Cu1	-161.09 (14)	O1—Gd1—O8—N4	97.0 (3)
O4—Gd1—O2—Cu1	18.9 (2)	N3—Gd1—O8—N4	-77.8 (5)
O1—Gd1—O2—Cu1	157.0 (2)	O10—N4—O9—Gd1	-171.4 (5)
N4—Gd1—O2—Cu1	95.71 (18)	O8—N4—O9—Gd1	7.1 (5)
N3—Gd1—O2—Cu1	-61.4 (2)	O2—Gd1—O9—N4	-38.4 (4)
O2—Cu1—O3—C19	-168.8 (4)	O3—Gd1—O9—N4	36.1 (3)
N2—Cu1—O3—C19	-2.0 (4)	O11—Gd1—O9—N4	-140.6 (3)
N1—Cu1—O3—C19	140.3 (5)	O6—Gd1—O9—N4	146.0 (3)
Gd1—Cu1—O3—C19	178.0 (4)	O8—Gd1—O9—N4	-4.2 (3)
O2—Cu1—O3—Gd1	13.20 (14)	O5—Gd1—O9—N4	122.3 (3)
N2—Cu1—O3—Gd1	-179.97 (16)	O12—Gd1—O9—N4	-148.8 (3)
N1—Cu1—O3—Gd1	-37.6 (6)	O4—Gd1—O9—N4	80.5 (3)
O2—Gd1—O3—C19	169.8 (4)	O1—Gd1—O9—N4	-77.5 (3)
O9—Gd1—O3—C19	43.1 (4)	N3—Gd1—O9—N4	135.5 (3)
O11—Gd1—O3—C19	-140.4 (3)	O13—N5—O11—Gd1	179.4 (5)
O6—Gd1—O3—C19	-58.5 (4)	O12—N5—O11—Gd1	-0.7 (5)
O8—Gd1—O3—C19	75.1 (3)	O2—Gd1—O11—N5	-142.4 (3)
O5—Gd1—O3—C19	-92.6 (4)	O3—Gd1—O11—N5	174.3 (3)
O4—Gd1—O3—C19	-4.2 (3)	O9—Gd1—O11—N5	-9.3 (3)
O1—Gd1—O3—C19	134.9 (3)	O6—Gd1—O11—N5	75.1 (3)
N4—Gd1—O3—C19	58.2 (3)	O8—Gd1—O11—N5	-73.2 (4)
N3—Gd1—O3—C19	-73.9 (4)	O5—Gd1—O11—N5	128.2 (3)
O2—Gd1—O3—Cu1	-12.16 (13)	O12—Gd1—O11—N5	0.4 (3)
O9—Gd1—O3—Cu1	-138.80 (14)	O4—Gd1—O11—N5	84.1 (3)
O11—Gd1—O3—Cu1	37.7 (2)	O1—Gd1—O11—N5	-77.3 (3)
O6—Gd1—O3—Cu1	119.6 (2)	N4—Gd1—O11—N5	-33.3 (4)
O8—Gd1—O3—Cu1	-106.83 (17)	N3—Gd1—O11—N5	101.1 (4)
O5—Gd1—O3—Cu1	85.5 (2)	O13—N5—O12—Gd1	-179.4 (5)
O4—Gd1—O3—Cu1	173.9 (2)	O11—N5—O12—Gd1	0.7 (4)
O1—Gd1—O3—Cu1	-47.02 (19)	O2—Gd1—O12—N5	42.6 (3)
N4—Gd1—O3—Cu1	-123.76 (16)	O9—Gd1—O12—N5	170.3 (3)
N3—Gd1—O3—Cu1	104.2 (3)	O11—Gd1—O12—N5	-0.4 (3)
O2—Gd1—O4—C18	-1.2 (4)	O6—Gd1—O12—N5	-86.9 (3)
O3—Gd1—O4—C18	5.2 (3)	O8—Gd1—O12—N5	141.1 (3)
O9—Gd1—O4—C18	-130.0 (3)	O5—Gd1—O12—N5	-52.2 (4)
O11—Gd1—O4—C18	118.9 (3)	O4—Gd1—O12—N5	-137.4 (3)
O6—Gd1—O4—C18	128.2 (4)	O1—Gd1—O12—N5	82.2 (3)
O8—Gd1—O4—C18	-75.1 (3)	N4—Gd1—O12—N5	157.2 (3)
O5—Gd1—O4—C18	76.2 (3)	N3—Gd1—O12—N5	-70.6 (4)

Fig. 1

